

Trees

Direct Penman-Monteith parameterizations for estimating stomatal conductance and modeling sap flow --Manuscript Draft--

Manuscript Number:	TSAF-D-16-00012R2	
Full Title:	Direct Penman-Monteith parameterizations for estimating stomatal conductance and modeling sap flow	
Article Type:	Original Article	
Keywords:	canopy conductance; Transpiration; trunk heat balance; Penman-Monteith; alpine forest	
Corresponding Author:	Josef Urban Mendel University in Brno CZECH REPUBLIC	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Mendel University in Brno	
Corresponding Author's Secondary Institution:		
First Author:	Jiri Kucera	
First Author Secondary Information:		
Order of Authors:	Jiri Kucera	
	Patricia Brito	
	Maria Soledad Jiemenez	
	Josef Urban	
Order of Authors Secondary Information:		
Funding Information:	Spanish Government (CGL2006-10210/BOS)	dr. Maria Soledad Jiemenez
	MSMT COST (LD 13017)	dr. Josef Urban
	Russian Academic Excellence Project 5-100	dr. Josef Urban
Abstract:	<p>The Penman-Monteith equation of evaporation is often combined with sap flow measurements to describe canopy transpiration and stomatal conductance. The traditional approach involves a two-step calculation. In the first step, stomatal conductance is computed using an inverted form of Penman-Monteith equation. The second step correlates these values with environmental factors. In this work, we present an improved approach for direct parameterization of the Penman-Monteith equation developed to compute diurnal courses of stand canopy conductance (g_c) from sap flow. The main advantages of this proposed approach versus using the classical approach are: (1) the calculation process is faster and involves fewer steps, (2) parameterization provides realistic values of canopy conductance, including conditions of low atmospheric vapor pressure deficit (D) whereas the traditional approach tends to yield unrealistic values for low D and (3) the new calculation method does not require enveloping curves to describe dependence of g_c on D and thus avoids subjective data selection but it still allows to visualize separable responses of g_c to environmental drivers (i.e. global radiation and vapor pressure deficit). The proposed approach was tested to calculate g_c and to model the sap flow of a high mountain <i>Pinus canariensis</i> forest. The new calculation method permitted us to describe the stand canopy conductance and stand sap flow in sub-hour resolution for both day and night conditions. Direct parameterization of the Penman-Monteith approach as implemented in this study proved sufficiently sensitive for detecting diurnal variation in g_c and for predicting sap flow from environmental variables under various atmospheric</p>	

evapotranspirative demands and differing levels of soil water availability.

Response to Reviewers:

Ulrich E. Luetttge
Editor-in-Chief
Trees - Structure and Function
Dear Prof. Dr. Luetttge

First of all we sincerely thank you for your editorial decision and for the useful comments and helpful suggestions of the reviewers. We now have considered and incorporated the reviewers' comments and suggestions in the revised text. Please find below our answers to the reviewers' specific comments. We are convinced that the revised manuscript has been significantly improved and will now be acceptable for publication in your Trees - Structure and Function.

Yours sincerely
Josef Urban, on behalf of the co-authors

Answers to the reviewers' specific comments:

Reviewer #3:

Authors' response: As suggested, we enlarged the introduction and the discussion sections. We explained the reasons for using the Jarvis and Lohammar models instead of the Ball Berry model. We much appreciate Ball Berry model (and its future modifications) and its use in the global circulation models and in the models which include net photosynthesis. We are aware of the limitations of the Lohammar and Jarvis-Stewart models that we use, and that is also why we proposed to include the parameter of minimal stomatal conductance, similar to the one used by the Ball Berry model. Comparing to the Lohammar model, the inclusion of net assimilation (and its correlation with the stomatal conductance) into the Ball Berry model reduced the number of free parameters, which is definitely a great advantage. On the other hand, when using the Ball Berry model, the net assimilation has to be either measured or simulated by the biochemical models. Many of the sap flow studies do not have data on photosynthesis available (even though the already published values from the similar sites or tree species can be used too). Also, unlike the sap flow, photosynthesis cannot be easily measured on the whole tree level which also brings along the scaling issues. Therefore the Jarvis-Stewart and Lohammar models are still used in the simple studies like ours for their simplicity, when the burden of measuring and modeling of photosynthesis does not justify the advantages of its inclusion. Moreover, the Ball Berry model has some limitations too. Even though the fixed relationship between the assimilation and stomatal conductance holds for the most of the situation, it changes sometimes. One example may be the drought (which occurred in our study and also in the study of Xu, Medvigy et al. (2016), where at Fig. 4a the sap flow almost do not change in the daytime between dry and wet season while the net assimilation decreases: maybe among others due to the changes in mesophyll conductance which affect the CO₂ transfer to the chloroplasts and do not such have effect on water vaporization) and the other example the extreme temperatures during heatwaves. We agree that there is a large array of the models which provide more mechanistic insight to the stomatal functioning than ours, among them the works of Xu, Medvigy et al. (2016) and Mirfenderesgi, Bohrer et al. (2016). On the other hand, there are even more detailed models of stomatal behavior, which include processes on the membranes of guard cells. All of them have their place in the tree research and none of them is ideal for all kind of applications. There are still papers being published which use the methodology we try to improve and that's why we think we have reason to care about this kind of models too. We included some more references of papers using approach similar to ours to highlight relevance of our study.

Reviewers' question: What is the physical rationale behind the use of the arctangent function in equation 4?

Authors' response: Some data suggested, that stomata response to D may not be always exponential and that stomata may slow down its opening at low vapor pressure deficit (i.e. Jones 2014, p. 140, Fig. 6.10, Bourne et al. 2015, Fig. 2), we introduced an arc tangential relationship allowing for more precise descriptions of stomata behavior at low D. Because gasometrical measurement suffer difficulties at large air humidities due to the condensations, there are not many data available and the proposed arctangential function may not be the best one. On the other hand, it yielded the highest R² and lowest AIC which suggest that its use may be rational.

Reviewers' question: The methodology of Ewers, Mackay et al. (2007) for estimating

stomatal conductance on the basis of observed sap flow does suffer from difficulties at low VPD, but in what scenarios would it be beneficial to back calculate sap flow and stomatal conductance from the PM model when neither are observed?

Authors' response: The analysis allows describing the response of g_c of the particular tree species to various environmental drivers (i.e. R_g or D). Sap flow calculated by this method may also be used for simulation of water use of the forests when it was not measured or to fill the gaps in the sap flow measurements.

Reviewers' question: Importantly, why would one chose to use this method to back calculate sap flow, over a mechanistic model that represents physical pathway of the soil-plant-atmosphere continuum as is becoming increasingly common using the models the authors cite in line 360 as well in the works of Xu, Medvigy et al. (2016) and Mirfenderesgi, Bohrer et al. (2016)?

Authors' response: There is a large array of the models to describe the tree water relations. Our method (in its 'classical' approach) is quite commonly used due to its simplicity and ability to describe separate responses of stomatal conductance to various environmental drivers. One advantage of this simple approach is that that almost everyone is able to calculate stomatal conductance. The other point is that in such a simple model there are not many parameters to be mutually adjusted. I admire the accuracy with which can the FETCH2 model predict the forest transpiration. On the other hand there may arise the questions about how accurately are the underlying processes described. For example in the work of Mirfenderesgi et al. (2016), Fig. 1b: is really the P50, the point where the 50 % of the xylem conductance is lost, in a pine as low as about the -1MPa? Or is it just a result of numerical solving and mutual adjustment of many parameters? Is the pine more vulnerable to cavitation than oak? How does it match to the published data? This is the point where the complex models can provide questions about plant and ecosystem functioning but we should be careful about taking the model results as answers.

Reviewers' question: L62-64: How can stomatal conductance be dependent on soil moisture and xylem architecture while excluding controls imposed through the actual xylem/tracheid pathway? I.e. soil moisture governs root water availability, which governs root water potential, which acts with xylem architecture to govern xylem water potential, which in turn governs leaf water potential which influences stomatal conductance.

Authors' response: We improved the wording at this point of the manuscript: L58 - 63

Reviewers' question: L68: Also importantly (Collatz, Ball et al. 1991) and (Ball, Woodrow et al. 1987)

Authors' response: We added these references to the text.

Reviewers' question: L88: Subject verb agreement 'substitutes'

Authors' response: Changed as suggested.

Reviewers' question: L89: The subject of this sentence is missing

Authors' response: The subject of the sentence was added.

Reviewers' question: L112: What is the site's typical VPD range since we are talking about why it is important not to have to discard data when VPD is low? What fraction of half hours would have to be discarded during a typical growing season because VPD is below 600kPa?

Authors' response: We included reference to the Fig. 2 and added the exact number of rejected values (818 out of the measured 1872 values) on the L255.

Reviewers' question: L140-Forward: This would be clearer if it were presented in a table for all variables used throughout the paper.

Authors' response: We created a table with abbreviations as suggested.

Reviewers' question: L170: What is the specific justification for the arctan term? How was this developed/determined?

Authors' response: We answered to this question in one of the previous questions of this letter and also added the explanation to the manuscript.

Reviewers' question: L174: Should be g_{max} ?

Authors' response: g_{lim} is the theoretical maximum of the arctangent function which may be higher than maximum g_s observed at zero D . Therefore we prefer to use the term g_{lim} .

Reviewers' question: L213-217: What were the soil water potential ranges during each period?

Authors' response: We added the SWP to the lines 215-222

Reviewers' question: L226: On what time scale, specifically?

Authors' response: We changed the wording to: 'These calculations were performed in 24 hour resolution within a period from 15 February to 1 September, 2008.'

Reviewers' question: L227-230 This makes no sense. Why would you parameterize a model for something only to change your parameters specifically when the model did not fit?

Authors' response: We tried to better explain as: 'In a two-week long period from 25 May to 10 June, following a rainfall event, gypsum block sensors indicated an increase in soil water potential which, however, was not entirely reflected in tree sap flow (Figs. 2b and 3). This prompted us to use linear interpolation for the SWP between the beginning and end of this period, from 25 May to 12 June and to use only these data for the model parameterization.'

Reviewers' question: L234-238: Did any or none of these include the effect of soil water potential?

Authors' response: None of them included the effect of SWP, we added this information on the lines 250-251.

Reviewers' question: L241-2: Were they not compared directly to field measurements of sap flow?

Authors' response: We improved the wording here: 'Values of modeled sap flow and canopy conductance were compared to the sap flow measured in the field and to a canopy conductance calculated from an inverted P-M equation (7).'

Reviewers' question: L247: True, but isn't your whole argument that your methodology is superior than the classic method particularly in cases of low D? In that case, why continue to not simulate times when D is low? Wouldn't the authors want to make the comparison during these times in particular?

Authors' response: We did two comparisons: First one for the period where classical approach work without problems to prove that proposed approach provide comparable results. Second comparison included also the period where classic method fails, to demonstrate that proposed method works even during this period. We also improved the text of the manuscript here.

Reviewers' question: L261: differences?

Authors' response: Changed as suggested

Reviewers' question: L264: discrepancies?

Authors' response: Changed as suggested.

Reviewers' question: L270: Most other stomatal conductance models include a g_{min} parameter (typically, g_o) which, again, raises the question why the authors chose to use the Lohammar model rather than a more broadly used and accepted model for stomatal conductance?

Authors' response: We believe that we explained the reasons at the beginning of this letter as well as in the introduction section of the manuscript.

Reviewers' question: L294: If this is the case, again, why not use a more broadly accepted g_s model that already includes a g_o term??

Authors' response: We believe that we explained the reasons at the beginning of this letter as well as in the introduction section of the manuscript.

Reviewers' question: L357-9: All of these models are for porous media flow. Could you provide examples of R-C models as well?

Authors' response: We added those models as suggested.

Reviewers' question: L360: The work of Matheny, Bohrer et al. (2015) and Oliva Carrasco, Bucci et al. (2015) have developed new measurement technologies for measuring water content directly in the stems of large trees at time steps smaller than a half hour.

Authors' response: We added these references and a one to the previous work on this topic to the text: L381-383.

Reviewers' question: L360-362: None of the models cited in L358 require the measurement of water potential to run or even to evaluate. All models solely require sap flux to parameterize the model in order to solve for water potentials, from which stomatal conductance is calculated.

Authors' response: We added a sentence of discussion on this topic on lines 376-379.

Reviewers' question: L390-1: Why is this not discussed in the results section?

Authors' response: We added these results to the lines 314-317.

Reviewers' question: Fig2: Soil water potential remained 100% constant from July through September?

Authors' response: -1.1 MPa was the limit of the equipment. This information is now on the lines L154 and 222.

Reviewers' question: Fig3: What do you mean by "modeled from wet period"? Is this using parameters that were established specifically for times during wet soils? Why is it more practical to re-parameterize a model in the middle of the growing season rather

than to include a more accurate representation of the dynamics that limit water availability?

Authors' response: We improved the figure caption. The explanation for the reason for modeling of the sap flow using the parameters from the period of unlimited water availability was included to the lines 271-272

[Click here to view linked References](#)

Direct Penman-Monteith parameterizations for estimating stomatal conductance and modeling sap flow

Jiří Kučera¹, Patricia Brito², María Soledad Jiménez², Josef Urban^{3, 4,*}

¹EMS Brno, Turistická 5, Brno, Czech Republic

²Departamento de Botánica, Ecología y Fisiología Vegetal, Universidad de la Laguna (ULL), Tenerife, Spain

³Department of Forest Botany, Dendrology and Geobiocenology, Mendel University in Brno, Czech Republic

⁴Siberian Federal University, Krasnoyarsk, Russia

*Corresponding author: josef.urban@email.cz, telephone: +420776041303

Abstract

The Penman-Monteith equation of evaporation is often combined with sap flow measurements to describe canopy transpiration and stomatal conductance. The traditional approach involves a two-step calculation. In the first step, stomatal conductance is computed using an inverted form of Penman-Monteith equation. The second step correlates these values with environmental factors. In this work, we present an improved approach for direct parameterization of the Penman-Monteith equation developed to compute diurnal courses of stand canopy conductance (g_c) from sap flow. The main advantages of this proposed approach versus using the classical approach are: (1) the calculation process is faster and involves fewer steps, (2) parameterization provides realistic values of canopy conductance, including conditions of low atmospheric vapor pressure deficit (D) whereas the traditional approach tends to yield unrealistic values for low D and (3) the new calculation method does not require enveloping curves to describe dependence of g_c on D and thus avoids subjective data selection but it still allows to visualize separable responses of g_c to environmental drivers (i.e. global radiation and vapor pressure deficit). The proposed approach was tested to calculate g_c and to model the sap flow of a high mountain *Pinus canariensis* forest. The new calculation method permitted us to describe the stand canopy conductance and stand sap flow in sub-hour resolution for both day and night conditions. Direct parameterization of the Penman-Monteith approach as implemented in this study proved sufficiently sensitive for detecting diurnal variation in g_c and for predicting sap flow from environmental variables

33 under various atmospheric evapotranspirative demands and differing levels of soil water
34 availability.

35

36 **Keywords:** canopy conductance; transpiration; trunk heat balance; Penman-Monteith;
37 alpine forest.

38

39 **Key message:** The novel approach for direct parameterization of the Penman-Monteith
40 equation was developed to compute diurnal courses of stand canopy conductance from sap
41 flow.

42 **1. Introduction**

43 In the average, transpiration accounts for 61 % of the evapotranspiration (Schlesinger
44 and Jasechko 2014). It means, that most of water evaporated from the ecosystems has to
45 pass through the plant and that its precise amount is regulated by the vegetation. Therefore,
46 the evaporation models have developed from the purely physical approach (i.e. Penman
47 1948) to the ones that implemented the principles of turbulent diffusion theory and plant
48 stomatal regulation (Monteith 1965). In coniferous and other stands, with strong coupling
49 between the canopy and the atmosphere, g_c (see Table 1 for the list of abbreviations) has a
50 major effect on transpiration (Jarvis and McNaughton 1986; Meinzer et al. 1997; Phillips and
51 Oren 1998; Hernandez-Santana et al. 2016). If evapotranspiration can be measured, actual
52 stomatal conductance can then be easily calculated by using an inverted Penman-Monteith
53 equation (Martin et al. 1997; Cienciala et al. 1997; Granier et al. 2007; Whitley et al. 2009;
54 Braun et al. 2010; Wang et al. 2014; Fu et al. 2016). This equation combined with sap flow
55 measurements provides a powerful tool for estimating canopy conductance for both whole-
56 tree and forest stand levels with results comparable to leaf level measurements with
57 subsequent upscaling (Ewers et al. 2007b).

58 The aim of calculating stomatal conductance with the P-M equation is to determine to
59 what extent g_c is dependent on environmental variables such as solar radiation and air vapor
60 pressure deficit (Lohammar et al. 1980; Oguntunde et al. 2007), which may be combined for
61 example with air temperature (Jarvis 1976; Stewart 1988; Sommer et al. 2002), soil moisture
62 (Stewart 1988; Granier and Loustau 1994; Harris et al. 2004), and eventually xylem
63 architecture (Zhang et al. 2011). A range of functions exist for these models, from a simple
64 linear approach (Oguntunde et al. 2007) using more complicated non-linear equations
65 relating two (Lohammar et al. 1980) or more (Jarvis 1976) environmental variables, to

66 complex models incorporating photosynthetic and biochemical regulation to measure
67 stomatal conductance (Ball et al. 1987; Collatz et al. 1991; Tardieu and Davies 1993;
68 Leuning et al. 1995; Jarvis and Davies 1998; Berry et al. 2010; Buckley et al. 2012;
69 Mirfenderesgi et al. 2016; Xu et al. 2016). However, parameterizing the lastly mentioned
70 models which include photosynthesis has become increasingly difficult due to varying
71 degrees of model sophistication and limited data availability for most field studies (Ward et
72 al., 2008). For this reason, empirical models are commonly selected to focus on two primary
73 environmental drivers: R_g and D . In conditions of limited soil water content, soil water deficit
74 may be added as a third driver.

75 There are two main types of the empirical (phenomenological) models of the stomatal
76 conductance. First group is represented by the Jarvis-Stewart (Stewart 1988) and the
77 Lohammar model (Lohammar et al. 1980). These models predict stomatal responses solely
78 on the base of environmental factors. This approach is still widely used because it is simple
79 and its modular structure makes it easy to incorporate in the larger models, and because its
80 responses to the environmental factors can be separated and visualized (Egea et al. 2011;
81 Buckley and Mott 2013). For its simplicity this kinds of models are also used to derive
82 stomatal conductances from the sap flow measurements (Lu et al. 1995; Cienicala et al.
83 1997; Whitehead 1998; Oren et al. 1999; Ewers and Oren 2000; Ewers et al. 2007a; García-
84 Santos et al. 2009; Wang et al. 2014; Fu et al. 2016). In such a cases, the simple models are
85 often preferred over the more recent Ball-Berry model and its modifications, which are
86 otherwise widely used in canopy and global circulation models (Verhoef and Egea 2014). The
87 main complication for the use of Ball-Berry model in derivation of g_c from the sap flow is that
88 it requires the knowledge of net assimilation. Net assimilation can be directly measured
89 (which data are not always available with sap flow measurements) and then plotted against
90 measured g_s or modeled from biochemistry of photosynthesis but it presents an extra effort
91 that does not need to be done when using the simpler Lohammar model. For the sake of
92 simplicity and to make our work comparable to similar studies (i.e. Oren et al. 1999; Ewers
93 and Oren 2000; Ewers et al. 2007a; García-Santos et al. 2009; Wang et al. 2014; Fu et al.
94 2016) we decided to use those kinds of empirical models that do not include the
95 photosynthesis.

96 In the classical process for deriving stomatal conductance from sap flow measurements,
97 g_c is calculated using the inverted form of the P-M equation followed by regression analyses
98 usually performed between individual environmental variables and g_s . One disadvantage of
99 this approach is that it requires using an enveloping curve for D and excluding data when D
100 rates are low to avoid unrealistically high values of g_s (Phillips and Oren 1998; Ewers and

101 Oren 2000; García-Santos et al. 2009). This paper proposes an improved way of modeling
102 canopy sap flow in a forest stand through direct parameterization of a P-M equation. Above
103 mentioned traditional approach first (i) calculates the g_c from inverted P-M equation, than (ii)
104 it rejects all data where g_c seem unrealistic (especially at low D), (iii) it searches for
105 dependence between g_c and weather conditions using regression analysis and suitable (i.e.
106 Jarvis-Stewart or Lohammar) model, (iv) substitutes that sub-model into the P-M equation
107 and calculates the sap flow. In contrast, proposed approach enables to perform modeling of
108 sap flow in one step and it does not require data rejection when D levels are low. The g_c and
109 stomatal response to R_g and D can be computed in a second step from the Lohammar
110 equation. We also propose a few modifications to the Lohammar equation. Because some
111 data suggested, that stomata response to D may not be always exponential (i.e. Jones 2014,
112 p. 140, Fig. 6.10, Bourne et al. 2015, Fig. 2), we introduced an arc tangential relationship
113 allowing for more precise descriptions of stomata behavior at low D. An additional parameter
114 describing minimal stomatal conductance (similarly to the parameters of 'b' or 'g₀' in Ball-
115 Berry model and its modifications) was then introduced permitting calculation of nocturnal
116 sap flow. The aim of this study is to improve how sap flow and stomatal conductance are
117 modelled from Penman-Monteith equation, to test the applicability of proposed approach for
118 modeling sap flow in a forest stand and to compare the results with those derived from using
119 the more traditional approach. The analysis allows describing the response of g_c of the
120 particular tree species to various environmental drivers (i.e. R_g or D). Sap flow calculated by
121 this method may also be used for simulation of water use of the forests when it was not
122 measured or to fill the gaps in the sap flow measurements.

123 **2. Material and methods**

124 **2.1. Site description**

125 The study was conducted in an open *P. canariensis* forest at its upper distribution limit
126 at 2,070 m a.s.l. in the Teide National Park (28°18'21.5"N, 16°34'5.8"W), Tenerife, Canary
127 Islands, Spain. At the time of the study (2008), the trees were 50–60 years old with a mean
128 canopy height of 10.3 m. The stand density was 291 trees ha⁻¹, with a basal area of 16.7 m²
129 ha⁻¹. The diameter at breast height (DBH) averaged 25.4 ± 9.3 cm (Fig. 1), and the plant
130 area index (LAI) was 3.6 (Brito et al. 2014). The field site is characterized by a
131 Mediterranean climate with alternating warm and dry periods from June to September, and a
132 cold and wet period from October to May. During the period 1971–1999, mean annual
133 precipitation recorded at the Portillo weather station located approximately 300 m southeast

134 of the study site was 368 mm, falling mostly during the winter months (December–
135 February; 190 mm), while summer was the driest season (June–August; 3 mm). The mean
136 annual air temperature was 10.7 °C. Daily means of global radiation, vapor pressure deficit,
137 soil water potential and daily sums of precipitation within the investigated period are at the
138 Fig. 2. The geologic substrate is of volcanic origin, and the soil at the study site is classified
139 as a leptosol.

140 **2.2. Field measurements**

141 Sap flow was measured from 25 January to 10 September 2008. Ten sample trees with
142 diameter at breast height (DBH) from 21 to 44 cm were selected for continuous sap flow
143 measurements. Tree sap flow was measured by the tissue heat balance method with
144 constant heating power (Čermák et al. 1973; Kučera et al. 1977) manufactured by EMS
145 Brno, Czech Republic (type P4.1, heating power 0.63 W). Electrode lengths of 70 and 80 mm
146 were used to cover the conductive sapwood depth. Thirty-minute averages from sap flow
147 values measured in one minute intervals were stored on a data logger. Scaling up of sap
148 flow from the sample trees to the stand level was based on the DBH of the sample trees and
149 diameter distribution of the trees in the forest stand (Čermák et al. 2004), as well as for the
150 study site, described in detail by Brito et al. (2014).

151 Environmental variables such as global radiation, air temperature, air humidity,
152 precipitation and wind speed were taken from a meteorological station (uEMSet 99 (EMS
153 Brno)) installed on an open plot close to the study site. Soil water potential between 0 and -
154 1.1 MPa, which was the limit of the equipment, was measured with three gypsum blocks at
155 depths of 25-30 cm (sensors GB2, Delmhorst, Inc., USA; datalogger ModuLog 1029, EMS
156 Brno, Czech Republic). All environmental data were recorded with a 30-minute resolution.

157 **2.3. Data processing**

158 2.3.1. New approach: derivation of canopy conductance by direct
159 parameterization of the Penman-Monteith equation

160 The analysis of canopy transpiration (E_p , mm h⁻¹) was made based on the P-M equation
161 with the simplification that for long periods, stand transpiration equals sap flow. The general
162 form of the P-M equation is:

$$163 \lambda E_p = \frac{\Delta(R_n - G) + \rho c_p D g_a}{\Delta + \gamma(1 + \frac{g_a}{g_c})} \quad (1)$$

164 and the aerodynamic conductance is:

$$165 \quad g_a = \frac{k^2 u}{\ln^2 \left[\frac{(z-z_d)/z_0}{z_0} \right]} \quad (2)$$

166 All above listed variables were directly measured or calculated. The only unknown variable
167 was canopy conductance g_c .

168 Parameterization of the P-M equation is based on the assumption that canopy
169 conductance g_c depends on solar radiation and D, according to a suitable formula. We used
170 a modified version of the Lohammar equation (Lohammar et al. 1980). The original form of
171 this equation supposes that stomatal opening is caused by solar radiation and that stomatal
172 closure results from high evaporative demands. Consequently, this proposes that stomatal
173 closure is complete at night. However, calculating transpiration response to night D values
174 called for an additional parameter (g_{\min}), similarly as in the Ball-Berry model (Ball et al.
175 1987). Some measured data suggested, that opening of stomata in response to the D may
176 not always follow strictly exponential manner (i.e. Jones 2014, p. 140, Fig. 6.10, Bourne et
177 al. 2015). Therefore, alongside with the exponential form of the model, we also applied
178 arctangent equation which gave us more flexibility when describing stomatal behavior under
179 conditions of low D. The original form of the Lohammar model was:

$$180 \quad g_c = \frac{R_g}{R_g + R_0} \cdot \frac{g_{\max}}{1 + a \cdot D} \quad (3)$$

181 and the proposed modification is:

$$182 \quad g_c = g_{\min} + \frac{R_g}{R_g + R_0} \cdot g_{\lim} \left(0.5 - \frac{1}{\pi} \arctg \left(\frac{D}{a} - b \right) \right) \quad (4)$$

184

185 The resulting form of the modified Penman-Monteith equation used for the parameterization
186 was:

$$187 \quad E = \frac{(\Delta \cdot (R_n - G) + \rho c_p D g_a) / \lambda}{\Delta + \gamma \cdot \left(1 + \frac{g_a}{g_{\min} + \frac{R_g + 1}{R_g + R_0} \cdot g_{\lim} \left(0.5 - \frac{1}{\pi} \arctg \left(\frac{D}{a} - b \right) \right)} \right)} \quad (5)$$

188 The parameters a , b , R_0 , g_{\min} and g_{\lim} were optimized by minimizing the residual sum of
189 squares of the observed and modeled conductances. Calculations were performed using non-
190 linear multivariate analysis with Mini32 statistical software (EMS Brno, Czech Republic).

191 Stomatal conductance (g_s) may be derived from canopy conductance (g_c) using the
192 following equation:

$$193 \quad g_c = g_s * LAI \quad (6)$$

194 The time lag between R_g , D (which determine foliar transpiration) and stem sap flow
195 were estimated performing time series cross-correlation analysis. The respective
196 environmental variable was shifted in thirty-minute step intervals and correlated to sap flow
197 until the highest R^2 was reached. Only the lagged environmental data were used for
198 parameterization of the P-M equation.

199 2.3.2. Classical approach

200 Results of the new approach were compared to a classically used method (Granier and
201 Loustau, 1994). Here, the canopy conductance was calculated from the inverted form of the
202 Penman-Monteith equation:

$$203 \quad \frac{1}{g_c} = \left[\frac{s}{\gamma} \left(\frac{R_g - \lambda E}{\lambda E} \right) - 1 \right] \frac{1}{g_a} + \frac{\rho_a c_p}{E \lambda \gamma} D \quad (7)$$

204 Subsequently, canopy conductance was modeled using both a modified Jarvis-Stewart
205 approach (Jarvis, 1976; Granier and Loustau, 1994) and a modified Lohammar model (eq. 4)
206 which was previously used for the direct parameterization of the P-M equation. The Jarvis-
207 Stewart model was used in the form:

$$208 \quad g_s = a \left(\frac{R_g}{R_g + b} \right) \left(\frac{1 - cD}{1 + dD} \right) + g_{min} \quad (8)$$

209 where a , b , c , d and g_{min} are estimated parameters based on the standard minimal sum of
210 squares criterion. Modeled canopy conductance was used for modeling the canopy sap flow
211 by applying the P-M approach. For parameterizations of equations 4 and 8 and classical
212 approach we used data when D was higher than 600 Pa (Ewers and Oren 2000). These
213 parameters estimated for high D were also use to model sap flow when the D was low.

214 2.3.3. Effect of soil water availability on sap flow

215 Three time periods characterized by contrasting soil water availability were apparent in
216 the study period. The first, characterized by non-limiting water availability ('wet' period;
217 mean soil water potential (SWP) was -0.02 MPa and it was always higher than -0.05 MPa
218 within this period), lasted from the beginning of the year to the end of April. The second, the
219 transient period, lasted from May to the middle of June. The SWP ranged between -0.05 and
220 -1.1 MPa most of the time. The last, dry period, occurred from the middle of June until the

221 end of the investigated period. The SWP in this period was lower than -1.1 MPa, which was
222 the lowest value that could be measured by the used equipment (Fig. 2). To avoid any
223 adverse effects of soil water availability on the stand transpiration, initial parameterization of
224 the proposed model was carried out for the period of non-limiting water access, from 27
225 February to 5 April, 2008. All comparisons between the proposed and classical approaches
226 were drawn from this same period.

227 We later modified eq. 4 in a way that enabled us to use the soil water potential data to
228 describe the effects of soil water availability on stomatal conductance:

$$229 \quad g_c = g_{\min} + \frac{R_g}{R_g + R_0} \cdot g_{\lim} \cdot \left(0.5 - \frac{1}{\pi} \arctg\left(\frac{D}{a} - b\right)\right) \cdot (c \cdot \ln(SWP) + d) \quad (9)$$

230 wherein SWP (MPa) represents the absolute value of soil water potential and c , d are
231 parameter estimates. These calculations were performed in 24 hour resolution within a
232 period from 15 February to 1 September, 2008. In a two-week long period from 25 May to
233 10 June, following a rainfall event, gypsum block sensors indicated an increase in soil water
234 potential which, however, was not entirely reflected in tree sap flow (Figs. 2b and 3). This
235 prompted us to use linear interpolation for the SWP between the beginning and end of this
236 period, from 25 May to 12 June and to use only these data for the model parameterization.

237 2.3.4. Comparison among the sub-models for dependence of stomatal 238 conductance on environmental variables and between classical and 239 proposed approach

240 Three kinds of models for canopy conductance dependence on environmental variables
241 (such as R_g , D) were applied during the process of direct parameterization. The Jarvis-
242 Stewart model (8), both the original and the modified Lohammar equation (3, 4), and the
243 three with the applied (or not applied) added parameter of minimal canopy conductance
244 (g_{\min}) – altogether provided six variants. These equations were included in the process of
245 direct parameterization of the P-M equation to calculate sap flow and canopy conductance.
246 Also, all these sub-models were used to calculate sap flow and g_c by classical approach.
247 Values of modeled sap flow and canopy conductance were compared to the sap flow
248 measured in the field and to a canopy conductance calculated from an inverted P-M equation
249 (7). All simulations intended for the comparison among the models were performed for the
250 period of non-limiting water access, from 27 February to 5 April, 2008 and therefore no
251 submodel for the soil water potential was applied. It is well known that the occurrence of low
252 D (high air humidity) leads to unrealistically high values of g_c calculated by the classical
253 approach. For this reason Ewers and Oren (2000) recommended to remove all data where

254 the D is lower than 600 Pa from the analysis. However, in our case this would mean
255 rejection of 818 out of the measured 1872 values. Therefore, we did two comparisons on
256 two different datasets. First one covered entire wet period and included all data, including
257 those when the D was lower than 600 Pa. The second dataset contained only 20 full days
258 when the D was higher than 600 Pa.

259 **3. Results**

260 **3.1. Measured sap flow**

261 Daily maximum levels of sap flow for individual sample trees were between 10 to 67 kg
262 day⁻¹ with maximal values of stand sap flow 0.89 mm day⁻¹ (Figure 3). Sap flow lagged
263 behind evapotranspiration demands of atmosphere, R_g and D. Estimated time lags between
264 the sap flow and climate variables were 60 minutes for R_g and 30 minutes for D.

265 **3.2. Modeling sap flow and stomatal conductance: direct** 266 **parameterization**

267 Our proposed model successfully predicted diurnal courses of stand sap flow during the
268 wet period, yielding $R^2 = 0.96$ and having the lowest AIC (Table 2), as soil moisture is not a
269 limiting factor during this part of the season. However, the model parameterization for the
270 wet season overestimated daily sums of sap flow during the transient and dry periods,
271 beginning in May (Fig. 3). Differences between measured and modeled sap flow may be
272 perceived as quantitative indicator of water deficit for the transpiration. The model provided
273 an acceptable description of tree sap flow on a diurnal basis (Figs. 4, 5), and even took into
274 account night flows. Discrepancies between measured and modeled values appeared after
275 rainfall, when it was apparent that the sap flow based model could not accurately predict
276 evaporation of intercepted water.

277 Parameterization of the P-M equation was further complicated by the occurrence of
278 night sap flow. The Lohammar equation, in its original form (3), accounts for the full closure
279 of the stomata and no cuticular transpiration, therefore for zero transpiration at night.
280 Introducing the parameter g_{\min} into the equation allowed us to estimate night flows. This
281 minimal nocturnal stomatal conductance (g_{\min}) was calculated as 0.00020 m s⁻¹. The highest
282 stand canopy conductance of 0.0033 m s⁻¹ and highest average conductance of 0.00107 m s⁻¹
283 were calculated during the wet period.

284 **3.3. Modeling sap flow and stomatal conductance: the classical**
285 **approach**

286 The classical approach, which applies an inverted P-M equation for calculating diurnal
287 courses of stomatal conductance with subsequent regression analysis between calculated g_c
288 and environmental variables (equations 7 and 8), also allowed for the estimation of stand
289 water use in a diurnal resolution. Calculations of canopy conductance upon low D were
290 subjected to an error typical for this method: when D was low, g_c unrealistically increased
291 (Fig. 4e). The error in estimating long-term water use (i.e. during the wet period) was similar
292 to the method of direct parameterization if only the periods of high D were considered.
293 Correlation between measured and modeled sap flow was slightly less than it was for the
294 model derived via direct parameterization (Table 2).

295 **3.4. Comparison of proposed and classical approach and of**
296 **different models for dependence of stomatal conductance on**
297 **environmental factors**

298 All models acceptably described diurnal courses of sap flow and stomatal conductance
299 when the D was high, with R^2 of linear regression between measured and predicted values
300 >0.94 and >0.87 for sap flow and stomatal conductance, respectively (Table 2). However,
301 when low D was included, only the proposed approach was able to predict the sap flow with
302 success. The classical approach yielded unrealistically high values of g_c (Fig. 4e) and there
303 was no correlation between g_c and environmental factors (R_g and D). Therefore we were not
304 able to model a sap flow when all data were used in a classical approach. A modified
305 Lohammar equation incorporating parameter of minimal stomatal conductance (g_{min}) proved
306 the most effective correlating with measured values, yielding highest R^2 and having the
307 lowest AIC of all models. Absolute values of sap flow and stomatal conductance were
308 generally more accurate when predicted by models incorporating parameter g_{min} , which were
309 able to account for the effect of night sap flow.

310 **3.5. Effect of soil water availability**

311 Sap flow dropped by 95 % during the dry period (Fig. 3). The transient period showed a
312 decrease in water use and lasted two months, from the end of April to the end of June. Daily
313 sums of sap flow were successfully predicted in all wet, transient and dry seasons.
314 Interpolation of SWP from 25 May to 12 June, after the single rain event during transient
315 period, helped to model the decline in sap flow. Otherwise the model would overestimate

316 measured sap flow and in this period would reach up to the values predicted from the
317 equation 5 (Fig. 3). The R^2 of a linear regression between measured and modeled sap flow
318 was 0.93.

319 **4. Discussion**

320 **4.1. Measured sap flow**

321 Low values of daily sums of stand sap flow reflected both harsh tree line conditions at
322 the altitude of roughly 2200 m a.s.l. (Fernández-Palacios and de Nicolás 1995) and low stand
323 density. Stand transpiration in one studied forest stand was lower than average at 0.80 mm
324 day^{-1} (with a maximum of 1.85 mm day^{-1}) in a lower situated stand of a *P. canariensis* (Luis
325 et al. 2005). Stomatal conductance was also lower than in previous studies for lower
326 altitudes (Wieser et al. 2002; Luis et al. 2005). A decrease in summer months of sap flow
327 and stomatal conductance, which occurred during the studied period, was typical only for the
328 higher altitudes, as summer drought at lower altitudes is often mitigated by a high frequency
329 of clouds that do not occur at high altitudes in the summer (Gieger and Leuschner 2004; Luis
330 et al. 2005; Wieser et al. 2006).

331 The estimated time lag of 1 hour between the diurnal course of global radiation and
332 stem sap flow, on the lower end of a commonly used range of 0 – 4 hours (Martin et al.
333 1997; Čermák et al. 2007; Whitley et al. 2009; Ward et al. 2013) was typical for the tree
334 species with reduced (more sparse) crowns and therefore smaller water storage capacity
335 (Anfodillo et al. 1998). Response to changes in D during the day were faster than to R_g as
336 once the pool of the easily accessible stored water is depleted, the hydraulic signals are
337 transmitted throughout the plant at a very high speed (Malone 1993). A lag of sap flow
338 compared to the potential evapotranspiration was also observed when the canopy was wet
339 from rain. Such a lag time may reach up to two hours (Langensiepen et al. 2009). In this
340 case, a simple P-M based model was not able distinguish between evaporation and
341 transpiration and therefore could not accurately predict sap flow.

342 **4.2. Comparison between direct parameterization and the classical** 343 **approach**

344 The classical approach for calculating g_c and transpiration (or sap flow) using the P-M
345 equation consists of several steps. In the first step, g_c is calculated from eq. 7. The second
346 step includes regression between R_g , D and g_c . Here, it is often necessary to introduce an

347 enveloping curve into the regression. Utilization of the inverted form of P-M equation, in
348 combination with the exponential equation characterizing the dependence of g_c on D , causes
349 an unrealistic increase of the g_c on low D (Fig. 4e). It happens because low sap flow is
350 divided by low D , which brings large error into the calculation. To avoid this phenomenon,
351 Ewers and Oren (2000) recommend excluding all data where D is lower than 600 Pa. This
352 often affects considerable parts of a dataset and in our case we would reject 44 % of the
353 data from the period of models comparison. The proposed approach using parameterization
354 of the P-M equation with simultaneous derivation of all parameters, using an arc tangential
355 relationship for D allowed us to predict g_c even on low D , and thus allowed us to predict sap
356 flow for any time of day. Furthermore, the one-step calculation process proved to be the
357 most consistent and resulted in the highest R^2 between observed and predicted sap flow and
358 in lowest AIC of the model (Table 2).

359 These two approaches have, however, common limitations. First of them is that they
360 are not able to distinguish between transpiration and evaporation. When the rain or dew
361 occurs intercepted water is evaporated first from the needle surfaces before the transpiration
362 follows. Evaporation consumes part of the radiation energy and increases the lag between
363 potential evapotranspiration and sap flow. Second issue relates to the hysteresis pattern
364 between sap flow and potential evapotranspiration (Oren et al. 2001; Matheny et al. 2014a;
365 Matheny et al. 2014b). Trees store considerable amount of water in the needles, branches,
366 stem and roots (Phillips et al. 2003; Warren et al. 2005; Čermák et al. 2007; Oliva Carrasco
367 et al. 2015; Urban et al. 2015; Mirfenderesgi et al. 2016). Resistances to the xylem water
368 transport make it easier to use some parts of the stored water for transpiration instead of
369 withdrawing it directly from soil (Sack and Holbrook 2006). Therefore, gradient of water
370 potential develops along the pathway of xylem water transport and time lag occurs between
371 transpiration and sap flow. Water stored in the tree is refilled during the periods of low
372 evapotranspiration demands. It brings further uncertainty into the modeling of sap flow from
373 weather data. One solution would be to introduce tree water potential or water storage into
374 the simulation, like in the resistance-capacitance circuit (Sperry et al. 1998; Steppe et al.
375 2006; Bonan et al. 2014) or porous media models (Bohrer et al. 2005; Chuang et al. 2006;
376 Janott et al. 2011; Mirfenderesgi et al. 2016). These models are able to numerically solve for
377 the water potential and they do not require it as a mandatory input variable because
378 continuous measurement of water potential in conifers is a significant challenge and these
379 data are often not available. Water potential can be estimated also indirectly, from the
380 dendrometers readings (Perämäki et al. 2001) or from changes in a stem water storage
381 measured by soil moisture sensors working on principle of time domain reflectometry (TDR)

382 (Irvine and Grace 1997) or frequency domain reflectometry (FDR) (Matheny et al. 2015;
383 Carrasco et al. 2015). Even though these indirect approaches have been developed, such
384 data are available only in a few studies. Therefore, with limited data availability,
385 parameterizing of the P-M equation is still a viable option to calculate g_c and to model stand
386 sap flow.

387 **4.3. Different models for dependence of g_c on environmental** 388 **variables**

389 Stomatal conductance models relate stomatal aperture to global radiation and vapor
390 pressure deficit. While R_g opens the stomata, D closes the stomata (Fig. 6). Strong
391 correlation between g_s and D lead to the assumption that a functional relationship exists
392 (Meinzer and Grantz 1990; Aphalo and Jarvis 1991; Addington et al. 2004). Both the Jarvis-
393 Stewart model incorporating the added parameter g_{min} (eq. 8) and the modified Lohammar
394 equation (eq. 4) acceptably described dependence of stomatal conductance on
395 environmental variables with only negligible differences in R^2 of the models within a
396 parameterized period.

397 The mechanism driving stomatal opening in the morning due to R_g led to an assumption
398 about stomatal movements only during daytime conditions (Shimazaki et al. 2007). However,
399 evidence of night sap flow response to ΔD suggests that large portions of water are lost
400 through cuticular transpiration (Burghardt and Riederer 2003), stomatal closure is incomplete
401 or that stomatal movements occur at night (Caird et al. 2007; Kavanagh et al. 2007; Zeppel
402 et al. 2011). Caird et al. (2007) summarized a large body of evidence showing that g_s
403 changes at night were based upon various driving factors, including D . We therefore
404 incorporated the parameter g_{min} characterizing night stomatal conductance into the eq. 4.
405 Inclusion of this parameter improved the correlation (Table 2). The parameter describing
406 minimal stomatal conductance is not a new idea. Similar parameter was used for example in
407 the Ball-Berry model. There is a difference between the g_{min} in our study and g_0 in the Ball-
408 Berry model. While g_{min} describes a true minimal stomatal conductance, g_0 is a stomatal
409 conductance at light compensation point of photosynthesis. Estimation of the g_{min} from the
410 sap flow data is further complicated by the changes in water content in plant tissues.
411 Therefore, the exact amount of water transpiration at night could not be accurately
412 described from the sap flux measurements as we were unable to separate plant refilling from
413 actual transpiration to the atmosphere (Wang et al. 2011). The hysteresis pattern observed
414 in the diurnal sap flux response to changing D (Oren et al. 2001) indicated that refilling
415 occurred. More complex models like FETCH2 (Mirfenderesgi et al. 2016) which recently

416 occurred would be needed to identify to what degree contributes aboveground tree water
417 storage to the tree water uptake and transpiration.

418 **4.4. Effect of soil moisture**

419 Soil water content is the other driving mechanism of stomatal aperture (Granier and
420 Loustau 1994; Zweifel et al. 2009; Brito et al. 2014). We defined a submodel describing the
421 effect of actual soil water potential on plant sap flow. This modification allowed for modeling
422 sap flow on a daily basis in wet, dry and transient periods of a season. However, a more
423 thorough understanding of the soil water potential in rooting zones (or of the predawn tree
424 water potential) and plant ecophysiological adaptation to imminent water stress is necessary.
425 In our case study, sap flow did not scale with measured soil water potential in a short (two-
426 week long) period after the rain event which occurred in the middle of the transient period,
427 when measured soil water potential increased from -0.5 MPa to almost zero, while sap flow
428 increased only slightly (Figs. 2, 3). Decoupling of the sap flow from measured soil water
429 potential may occur for two reasons. The first may be that it is linked to the position of soil
430 water sensors, which may not be able to describe true water availability for the tree root
431 system (as rooting depth can easily reach 15 m, while the sensors were installed in 30 cm
432 depth (Luis et al. 2005; Brito et al. 2015)). The second relates to a tree's ecophysiological
433 adaptation to water stress (i.e. needle shedding, which was observed at the site (BrITO et al.
434 2014), an accumulation of ABA, or embolism of xylem tracheids). Therefore, it is advisable to
435 monitor predawn tree water potential, as an indicator of a tree water status, and to monitor
436 longer-term tree adaptation to water stress.

437 **5. Conclusions**

- 438 1. Direct parameterization of the Penman-Monteith equation with an incorporated
439 modified Lohammar model is a suitable approach to derive diurnal courses of
440 stomatal conductance of the forest stand. Closer correlation between measured
441 and calculated sap flow was found using the proposed approach by applying the
442 direct parameterization than was observed using the classical calculation method.
- 443 2. Direct parameterization of the P-M equation allowed for calculation of canopy
444 conductance upon low D and avoided unrealistic increases of computed g_c , typical
445 for the classical approach of calculation. Furthermore, the direct parameterization
446 approach does not require computing the enveloping curve and thus avoids
447 subjective selection of the data by the user. Moreover, direct parameterization

448 allows for calculation of g_c and modeling sap flow to occur in one step. This makes
449 it possible to quickly and automatically repeat the calculation, i.e. for different time
450 periods.

451 3. The proposed modification of the Lohammar et al. (1980) equation successfully
452 described dependence of canopy conductance on measured environmental
453 variables (global radiation and vapor pressure deficit). Its modified form allowed for
454 more precise descriptions of g_c , especially upon low D , then the commonly used
455 three-parameter function. Including a parameter describing minimal stomatal
456 aperture allowed for modeling of night sap flow.

457 **Author Contribution Statement**

458 JK contributed to the work by devising of the methodology, data analysis and work on the
459 manuscript. MJ and PB performed the field work and contributed to the writing of the
460 manuscript. JU contributed to the data analysis and writing of the manuscript.

461 **Funding**

462 This work was supported by EMS Brno, the Spanish Government (CGL2006-10210/BOS co-
463 financed by FEDER, CGL2010-21366-C04-04 MCI), Russian Academic Excellence Project 5-
464 100 and the Czech project MSMT COST LD 13017 under the framework of the COST FP1106
465 network STReESS. P.B. received a fellowship from Canarian Agency for Research, Innovation
466 and Information Society (ACIISI) co-financed by FEDER.

467 **Acknowledgement**

468 The authors express their gratitude to National Park's Network for permission to work in
469 Teide National Park. We wish to thank A. Orsini for her work on language editing of the
470 manuscript.

471 **Conflict of Interest**

472 The authors declare that they have no conflict of interest.

References

- 473 Addington R, Mitchell R, Oren R, Donovan L (2004) Stomatal sensitivity to vapor pressure
474 deficit and its relationship to hydraulic conductance in *Pinus palustris*. *Tree Physiol*
475 24:561–569.
- 476 Anfodillo T, Rento S, Carraro V, et al. (1998) Tree water relations and climatic variations at

477 the alpine timberline: seasonal changes of sap flux and xylem water potential in *Larix*
478 *decidua* Miller, *Picea abies* (L.) Karst. and *Pinus cembra* L. *Ann For Sci* 55:159–172.

479 Aphalo P, Jarvis P (1991) Do stomata respond to relative humidity? *Plant, Cell Environ*
480 14:127–132.

481 Ball J, Woodrow I, Berry J (1987) A model predicting stomatal conductance and its
482 contribution to the control of photosynthesis under different environmental conditions.
483 *Prog Photosynth Res* IV:221–224.

484 Berry JA, Beerling DJ, Franks PJ (2010) Stomata: key players in the earth system, past and
485 present. *Curr Opin Plant Biol* 13:233–240. doi: 10.1016/j.pbi.2010.04.013

486 Bohrer G, Mourad H, Laursen TA, et al. (2005) Finite element tree crown hydrodynamics
487 model (FETCH) using porous media flow within branching elements: A new
488 representation of tree hydrodynamics. *Water Resour Res* 41:1–17. doi:
489 10.1029/2005WR004181

490 Bonan GB, Williams M, Fisher RA, Oleson KW (2014) Modeling stomatal conductance in the
491 earth system: Linking leaf water-use efficiency and water transport along the soil-plant-
492 atmosphere continuum. *Geosci Model Dev* 7:2193–2222. doi: 10.5194/gmd-7-2193-
493 2014

494 Bourne AE, Haigh AM, Ellsworth DS (2015) Stomatal sensitivity to vapour pressure deficit
495 relates to climate of origin in *Eucalyptus* species. *Tree Physiol* 35:266–278. doi:
496 10.1093/treephys/tpv014

497 Braun S, Schindler C, Leuzinger S (2010) Use of sap flow measurements to validate stomatal
498 functions for mature beech (*Fagus sylvatica*) in view of ozone uptake calculations.
499 *Environ Pollut* 158:2954–2963. doi: 10.1016/j.envpol.2010.05.028

500 Brito P, Lorenzo JR, González-rodríguez ÁM, et al. (2015) Canopy transpiration of a semi arid
501 *Pinus canariensis* forest at a treeline ecotone in two hydrologically contrasting years.
502 *Agric For Meteorol* 201:120–127. doi: 10.1016/j.agrformet.2014.11.008

503 Brito P, Lorenzo JR, González-Rodríguez ÁM, et al. (2014) Canopy transpiration of a *Pinus*
504 *canariensis* forest at the tree line: implications for its distribution under predicted
505 climate warming. *Eur J For Res* 133:491–500. doi: 10.1007/s10342-014-0779-5

506 Buckley TN, Mott KA (2013) Modelling stomatal conductance in response to environmental
507 factors. *Plant Cell Environ* 36:1691–1699. doi: 10.1111/pce.12140

508 Buckley TN, Turnbull TL, Adams MA (2012) Simple models for stomatal conductance derived
509 from a process model: cross-validation against sap flux data. *Plant, Cell Environ*
510 35:1647–1662. doi: 10.1111/j.1365-3040.2012.02515.x

511 Burghardt M, Riederer M (2003) Ecophysiological relevance of cuticular transpiration of
512 deciduous and evergreen plants in relation to stomatal closure and leaf water potential.
513 *J Exp Bot* 54:1941–1949. doi: 10.1093/jxb/erg195

514 Caird M, Richards J, Donovan L (2007) Nighttime stomatal conductance and transpiration in
515 C3 and C4 plants. *Plant Physiol* 143:4–10. doi: 10.1104/pp.106.092940

516 Carrasco LO, Bucci SJ, Di Francescantonio D, et al. (2015) Water storage dynamics in the
517 main stem of subtropical tree species differing in wood density, growth rate and life
518 history traits. *Tree Physiol* 35:354–365. doi: 10.1093/treephys/tpu087

519 Čermák J, Deml M, Penka M (1973) A new method of sap flow rate determination in trees.
520 *Biol Plant* 15:171–178.

521 Čermák J, Kučera J, Bauerle WL, et al. (2007) Tree water storage and its diurnal dynamics
522 related to sap flow and changes in stem volume in old-growth Douglas-fir trees. *Tree*

- 523 Physiol 27:181–198.
- 524 Čermák J, Kučera J, Nadezhdina N (2004) Sap flow measurements with some
525 thermodynamic methods, flow integration within trees and scaling up from sample trees
526 to entire forest stands. *Trees* 18:529–546. doi: 10.1007/s00468-004-0339-6
- 527 Chuang Y, Oren R, Bertozzi AL, et al. (2006) The porous media model for the hydraulic
528 system of a conifer tree: Linking sap flux data to transpiration rate. *Ecol Modell*
529 191:447–468. doi: 10.1016/j.ecolmodel.2005.03.027
- 530 Cienciala E, Kučera J, Lindroth A, et al. (1997) Canopy transpiration from a boreal forest in
531 Sweden during a dry year. *Agric For Meteorol* 86:157–167. doi: 10.1016/S0168-
532 1923(97)00026-9
- 533 Collatz GJ, Ball JT, Grivet C, Berry JA (1991) Physiological and environmental regulation of
534 stomatal conductance, photosynthesis and transpiration: a model that includes a
535 laminar boundary layer. *Agric For Meteorol* 54:107–136. doi: 10.1016/0168-
536 1923(91)90002-8
- 537 Egea G, Verhoef A, Vidale PL (2011) Towards an improved and more flexible representation
538 of water stress in coupled photosynthesis-stomatal conductance models. *Agric For*
539 *Meteorol* 151:1370–1384. doi: 10.1016/j.agrformet.2011.05.019
- 540 Ewers BE, Mackay DS, Samanta S (2007a) Interannual consistency in canopy stomatal
541 conductance control of leaf water potential across seven tree species. *Tree Physiol*
542 27:11–24. doi: 10.1093/treephys/27.1.11
- 543 Ewers BE, Oren R (2000) Analyses of assumptions and errors in the calculation of stomatal
544 conductance from sap flux measurements. *Tree Physiol* 20:579–589.
- 545 Ewers BE, Oren R, Kim H, et al. (2007b) Effects of hydraulic architecture and spatial variation
546 in light on mean stomatal conductance of tree branches. *Plant, Cell Environ* 30:483–
547 496. doi: 10.1111/j.1365-3040.2007.01636.x
- 548 Fernández-Palacios J, de Nicolás J (1995) Altitudinal pattern of vegetation variation on
549 tenerife. *J Veg Sci* 6:183–190.
- 550 Fu S, Sun L, Luo Y (2016) Combining sap flow measurements and modelling to assess water
551 needs in an oasis farmland shelterbelt of *Populus simonii* Carr in Northwest China. *Agric*
552 *Water Manag* 177:172–180. doi: 10.1016/j.agwat.2016.07.015
- 553 García-Santos G, Bruijnzeel LA, Dolman AJ (2009) Modelling canopy conductance under wet
554 and dry conditions in a subtropical cloud forest. *Agric For Meteorol* 149:1565–1572. doi:
555 10.1016/j.agrformet.2009.03.008
- 556 Gieger T, Leuschner C (2004) Altitudinal change in needle water relations of *Pinus*
557 *canariensis* and possible evidence of a drought-induced alpine timberline on Mt. Teide,
558 Tenerife. *Flora* 109:100–109.
- 559 Granier A, Loustau D (1994) Measuring and modelling the transpiration of a maritime pine
560 canopy from sap-flow data. *Agric For Meteorol* 71:61–81.
- 561 Granier A, Reichstein M, Bréda N, et al. (2007) Evidence for soil water control on carbon and
562 water dynamics in European forests during the extremely dry year: 2003. *Agric For*
563 *Meteorol* 143:123–145.
- 564 Harris P, Huntingford C, Cox PM, et al. (2004) Effect of soil moisture on canopy conductance
565 of Amazonian rainforest. *Agric For Meteorol* 122:215–227. doi:
566 10.1016/j.agrformet.2003.09.006
- 567 Hernandez-Santana V, Fernández J, Rodriguez-Dominguez C, et al. (2016) The dynamics of
568 radial sap flux density reflects changes in stomatal conductance in response to soil and

- 569 air water deficit. *Agric For Meteorol* 219:92–101.
- 570 Irvine J, Grace J (1997) Continuous measurements of water tensions in the xylem of trees
571 based on the elastic properties of wood.
- 572 Janott M, Gayler S, Gessler A, et al. (2011) A one-dimensional model of water flow in soil-
573 plant systems based on plant architecture. *Plant Soil* 341:233–256. doi:
574 10.1007/s11104-010-0639-0
- 575 Jarvis AJ, Davies WJ (1998) The coupled response of stomatal conductance to
576 photosynthesis and transpiration. *J Exp Bot* 49:399–406.
- 577 Jarvis P, McNaughton K (1986) Stomatal control of transpiration: scaling up from leaf to
578 region. *Adv Ecol Res* 15:1–49.
- 579 Jarvis PG (1976) The Interpretation of the Variations in Leaf Water Potential and Stomatal
580 Conductance Found in Canopies in the Field. *Philos Trans R Soc B Biol Sci* 273:593–610.
581 doi: 10.1098/rstb.1976.0035
- 582 Jones HG (2014) *Plants and Microclimate. A Quantitative Approach to Environmental Plant*
583 *Physiology*. doi: 10.1017/CBO9781107415324.004
- 584 Kavanagh KL, Pangle R, Schotzko AD (2007) Nocturnal transpiration causing disequilibrium
585 between soil and stem predawn water potential in mixed conifer forests of Idaho. *Tree*
586 *Physiol* 27:621–9.
- 587 Kučera J, Čermák J, Penka M (1977) Improved thermal method of continual recording the
588 transpiration flow rate dynamics. *Biol Plant* 19:413–420.
- 589 Langensiepen M, Fuchs M, Bergamaschi H, et al. (2009) Quantifying the uncertainties of
590 transpiration calculations with the Penman–Monteith equation under different climate
591 and optimum water supply conditions. *Agric For Meteorol* 149:1063–1072. doi:
592 10.1016/j.agrformet.2009.01.001
- 593 Leuning R, Kelliher FM, Pury DGG De, Schulze ED (1995) Leaf nitrogen, photosynthesis,
594 conductance and transpiration: scaling from leaves to canopies. *Plant Cell Environ*
595 18:1183–1200.
- 596 Lohammar T, Larsson S, Linder S, Falk S (1980) FAST: Simulation Models of Gaseous
597 Exchange in Scots Pine. *Ecol Bull* 32:505–523.
- 598 Lu P, Biron P, Bréda N, Granier A (1995) Water relations of adult Norway spruce (*Picea abies*
599 (L) Karst) under soil drought in the Vosges mountains: water potential, stomatal
600 conductance and transpiration. *Ann des Sci For* 52:117–129. doi:
601 10.1051/forest:19950203
- 602 Luis V, Jimenez M, Morales D, et al. (2005) Canopy transpiration of a Canary Islands pine
603 forest. *Agric For Meteorol* 135:117–123. doi: 10.1016/j.agrformet.2005.11.009
- 604 Malone M (1993) Hydraulic signals. *Philos Trans R Soc London, Biol Sci* 341:33–39.
- 605 Martin T, Brown K, Cermak J, et al. (1997) Crown conductance and tree and stand
606 transpiration in a second-growth *Abies amabilis* forest. *Can J For Res* 27:797–808.
- 607 Matheny A, Bohrer G, Stoy P, et al. (2014a) Characterizing the diurnal patterns of errors in
608 the prediction of evapotranspiration by several land-surface models: An NACP analysis.
609 *J Geophys Res* 119:1458–1473. doi: 10.1002/2014JG002623.Received
- 610 Matheny A, Bohrer G, Vogel C, et al. (2014b) Species-specific transpiration responses to
611 intermediate disturbance in a northern hardwood forest. *J Geophys Res Biogeosciences*
612 119:1–20. doi: 10.1002/2014JG002804.Received
- 613 Matheny AM, Bohrer G, Garrity SR, et al. (2015) Observations of stem water storage in trees

- 614 of opposing hydraulic strategies. *Ecosphere* 6:1–13. doi: 10.1890/ES15-00170.1
- 615 Meinzer FC, Andrade JL, Goldstein G, et al. (1997) Control of transpiration from the upper
616 canopy of a tropical forest: the role of stomatal, boundary layer and hydraulic
617 architecture components. *Plant, Cell Environ* 20:1242–1252. doi: 10.1046/j.1365-
618 3040.1997.d01-26.x
- 619 Meinzer FC, Grantz DA (1990) Stomatal and hydraulic conductance in growing sugarcane-
620 stomatal adjustment to water transport capacity. *Plant Cell Environ* 13:383–388.
- 621 Mirfenderesgi G, Bohrer G, Matheny AM, et al. (2016) Tree-level hydrodynamic approach for
622 modeling aboveground water storage and stomatal conductance illuminates the effects
623 of tree hydraulic strategy. *J Geophys Res Biogeosciences* 121:1792–1813. doi:
624 10.1002/2016JG003467
- 625 Monteith JL (1965) Evaporation and environment. *Symp Soc Exp Biol* 19:205–234.
- 626 Oguntunde P, Vandegiesen N, Savenije H (2007) Measurement and modelling of
627 transpiration of a rain-fed citrus orchard under subhumid tropical conditions. *Agric*
628 *Water Manag* 87:200–208. doi: 10.1016/j.agwat.2006.06.019
- 629 Oliva Carrasco L, Bucci SJ, Di Francescantonio D, et al. (2015) Water storage dynamics in the
630 main stem of subtropical tree species differing in wood density, growth rate and life
631 history traits. *Tree Physiol* 35:354–365. doi: 10.1093/treephys/tpu087
- 632 Oren R, Sperry J, Ewers B, et al. (2001) Sensitivity of mean canopy stomatal conductance to
633 vapor pressure deficit in a flooded *Taxodium distichum* L. forest: hydraulic and non-
634 hydraulic effects. *Oecologia* 126:21–29. doi: 10.1007/s004420000497
- 635 Oren R, Sperry JS, Katul GG, et al. (1999) Survey and synthesis of intra- and interspecific
636 variation in stomatal sensitivity to vapour pressure deficit. *Plant, Cell Environ* 22:1515–
637 1526. doi: 10.1046/j.1365-3040.1999.00513.x
- 638 Penman HL (1948) Natural evaporation from open water, bare soil and grass. *Proc R Soc*
639 *Lond A Math Phys Sci* 193:120–145.
- 640 Perämäki M, Nikinmaa E, Sevanto S, et al. (2001) Tree stem diameter variations and
641 transpiration in Scots pine: an analysis using a dynamic sap flow model. *Tree Physiol*
642 21:889–97.
- 643 Phillips N, Oren R (1998) A comparison of daily representations of canopy conductance
644 based on two conditional time-averaging methods and the dependence of daily
645 conductance on environmental factors. *Ann des Sci For* 55:217–235. doi:
646 10.1051/forest:19980113
- 647 Phillips N, Ryan M, Bond B, et al. (2003) Reliance on stored water increases with tree size in
648 three species in the Pacific Northwest. *Tree Physiol* 23:237–245.
- 649 Sack L, Holbrook NM (2006) Leaf hydraulics. *Annu Rev Plant Biol* 57:361–81. doi:
650 10.1146/annurev.arplant.56.032604.144141
- 651 Schlesinger WH, Jasechko S (2014) Transpiration in the global water cycle. *Agric For*
652 *Meteorol* 189–190:115–117. doi: 10.1016/j.agrformet.2014.01.011
- 653 Shimazaki K, Doi M, Assmann SM, Kinoshita T (2007) Light regulation of stomatal movement.
654 *Annu Rev Plant Biol* 58:219–47. doi: 10.1146/annurev.arplant.57.032905.105434
- 655 Sommer R, Sáb TDDA, Vielhauera K, et al. (2002) Transpiration and canopy conductance of
656 secondary vegetation in the eastern Amazon. *Agric For Meteorol* 112:103–121. doi:
657 10.1016/S0168-1923(02)00044-8
- 658 Sperry JS, Adler FR, Campbell GS, Comstock JP (1998) Limitation of plant water use by
659 rhizosphere and xylem conductance: Results from a model. *Plant, Cell Environ* 21:347–

- 660 359. doi: 10.1046/j.1365-3040.1998.00287.x
- 661 Steppe K, De Pauw DJW, Lemeur R, Vanrolleghem P a (2006) A mathematical model linking
662 tree sap flow dynamics to daily stem diameter fluctuations and radial stem growth. *Tree*
663 *Physiol* 26:257–273. doi: 10.1093/treephys/26.3.257
- 664 Stewart J (1988) Modelling surface conductance of pine forest. *Agric For Meteorol* 43:19–35.
- 665 Tardieu F, Davies WJ (1993) Integration of hydraulic and chemical signalling in the control of
666 stomatal conductance and water status of droughted plants. *Plant Cell Environ* 16:341–
667 349.
- 668 Urban J, Čermák J, Ceulemans R (2015) Above- and below-ground biomass, surface and
669 volume, and stored water in a mature Scots pine stand. *Eur J For Res* 134:61–74. doi:
670 10.1007/s10342-014-0833-3
- 671 Verhoef A, Egea G (2014) Modeling plant transpiration under limited soil water: Comparison
672 of different plant and soil hydraulic parameterizations and preliminary implications for
673 their use in land surface models. *Agric For Meteorol* 191:22–32. doi:
674 10.1016/j.agrformet.2014.02.009
- 675 Wang H, Guan H, Deng Z, Simmons CT (2014) Optimization of canopy conductance models
676 from concurrent measurements of sap flow and stem water potential on Drooping
677 Sheoak in South Australia. *Water Resour Res* 50:6154–6167. doi:
678 10.1002/2013WR014818
- 679 Wang H, Zhao P, Holscher D, et al. (2011) Nighttime sap flow of *Acacia mangium* and its
680 implications for nighttime transpiration and stem water storage. *J Plant Ecol* 2011. doi:
681 10.1093/jpe/rtr025
- 682 Ward EJ, Bell DM, Clark JS, Oren R (2013) Hydraulic time constants for transpiration of
683 loblolly pine at a free-air carbon dioxide enrichment site. *Tree Physiol* 33:123–134. doi:
684 10.1093/treephys/tps114
- 685 Ward EJ, Oren R, Sigurdsson BD, et al. (2008) Fertilization effects on mean stomatal
686 conductance are mediated through changes in the hydraulic attributes of mature
687 Norway spruce trees. *Tree Physiol* 28:579–596.
- 688 Warren JM, Meinzer FC, Brooks JR, Domec JC (2005) Vertical stratification of soil water
689 storage and release dynamics in Pacific Northwest coniferous forests. *Agric For Meteorol*
690 130:39–58. doi: 10.1016/j.agrformet.2005.01.004
- 691 Whitehead D (1998) Regulation of stomatal conductance and transpiration in forest
692 canopies. *Tree Physiol* 18:633–644.
- 693 Whitley R, Medlyn B, Zeppel M, et al. (2009) Comparing the Penman–Monteith equation and
694 a modified Jarvis–Stewart model with an artificial neural network to estimate stand-
695 scale transpiration and canopy conductance. *J Hydrol* 373:256–266. doi:
696 10.1016/j.jhydrol.2009.04.036
- 697 Wieser G, Luis VC, Cuevas E (2006) Quantification of ozone uptake at the stand level in a
698 *Pinus canariensis* forest in Tenerife, Canary Islands: An approach based on sap flow
699 measurements. *Environ Pollut* 140:383–386. doi: 10.1016/j.envpol.2005.12.003
- 700 Wieser G, Peters J, Luis VC, et al. (2002) Ecophysiological Studies on the Water Relations in
701 a *Pinus canariensis* Stand, Tenerife, Canary Islands. *Phyton (B Aires)* 42:291–304.
- 702 Xu X, Medvigy D, Powers JS, et al. (2016) Diversity in plant hydraulic traits explains seasonal
703 and inter-annual variations of vegetation dynamics in seasonally dry tropical forests.
704 *New Phytol* 212:80–95. doi: 10.1111/nph.14009
- 705 Zeppel MJB, Lewis JD, Chaszar B, et al. (2011) Nocturnal stomatal conductance responses to

706 rising [CO₂], temperature and drought. *New Phytol* no-no. doi: 10.1111/j.1469-
707 8137.2011.03993.x

708 Zhang Y, Oren R, Kang S (2011) Spatiotemporal variation of crown-scale stomatal
709 conductance in an arid *Vitis vinifera* L. cv. Merlot vineyard: direct effects of hydraulic
710 properties and indirect effects of canopy leaf area. *Tree Physiol* 1–18. doi:
711 10.1093/treephys/tpr120

712 Zweifel R, Rigling A, Dobbertin M (2009) Species-specific stomatal response of trees to
713 drought - a link to vegetation dynamics? *J Veg Sci* 20:442–454. doi: 10.1111/j.1654-
714 1103.2009.05701.x

715

716 Tables

717 Table 1. A summary of abbreviations and symbols used in the equations and throughout the
718 text.

a, b, c, d	parameters
AIC	Akaike's information criterion
c_p	specific heat of air [J m^{-3}]
D	vapor pressure deficit [Pa]
G	soil heat flux [W m^{-2}]
g_a	aerodynamic conductance [m s^{-1}]
g_c	canopy conductance [m s^{-1}]
g_{lim}	parameter related to maximal canopy conductance [m s^{-1}]
g_{min}	parameter of minimal canopy conductance [m s^{-1}]
g_s	stomatal conductance [m s^{-1}]
k	von Karman constant [-]
LAI	leaf area index [-]
P-M equation	Penman-Monteith equation
R_0	half light saturation [W m^{-2}]
R_g	global radiation [W m^{-2}]
R_n	net radiation [W m^{-2}]
SWP	soil water potential [MPa]
u	wind speed [m s^{-1}]
z	wind speed measurement height [m]
z_d	zero plane displacement [m]
z_o	canopy roughness [m]
γ	psychrometric constant [Pa K^{-1}]
Δ	slope of saturation water vapor pressure deficit [Pa K^{-1}]
λ	water heat capacity [J kg^{-1}]
ρ	density of dry air [kg m^{-3}]

719

720

721 Table 2. Index for determining of linear regression (R^2) between measured sap flow (or
 722 canopy conductance) and those simulated by the model. Akaike's information criterion
 723 (AIC) for various models. Simulated values of sap flow and stomatal conductance comes
 724 from following models: the Jarvis-Stewart model (eq. 8), the modified Lohammar model
 725 (eq. 4), the original Lohammar model (eq. 3) and values of sap flow and canopy
 726 conductance calculated from an inverted P-M equation (eq. 7). Models incorporating
 727 additional parameters of minimal stomatal conductance are labeled ' g_{min} '.
 728

		D > 600 Pa		Entire period			
		R^2	AIC	R^2	AIC		
Transpiration	Proposed model	Jarvis-Stewart g_{min}	0.955	-4503	0.934	-7128	
		Jarvis-Stewart original	0.953	-4440	0.932	-7052	
		Lohammar modified	0.954	-4447	0.933	-7058	
		Lohammar modified g_{min}	0.956	-4515	0.935	-7139	
		Lohammar original	0.949	-4417	0.929	-7022	
		Lohammar original g_{min}	0.950	-4464	0.929	-7084	
		Classical model	Lohammar modified	0.946	-4445	NA	NA
	Jarvis-Stewart g_{min}	0.940	-4412	NA	NA		
	Canopy conductance	Proposed model	Jarvis-Stewart g_{min}	0.907	-7080	0.000	
			Jarvis-Stewart original	0.907	-6999	0.000	
Lohammar modified			0.908	-7005	0.000		
Lohammar modified g_{min}			0.909	-7092	0.000		
Lohammar original			0.904	-6985	0.000		
Lohammar original g_{min}			0.903	-7058	0.000		
Classical model			Lohammar modified	0.887	-7028	0.000	
Jarvis-Stewart g_{min}		0.879	-7011	0.000			

729

Figure captions

Fig. 1. Diameter distribution of the pine trees at the research site, their number per ha (black bars) and the number and distribution of the sample trees used for sap flow measurements (grey bars).

Fig. 2. Daily means of global radiation (R_g), vapor pressure deficit (D, panel A), daily means of soil water potential (SWP) and daily sums of precipitation (panel B). Wet and dry periods are highlighted with a grey background. Dotted line on the panel B indicates values of SWP used to parametrize the model (9) after the rain event in the transient period.

Fig. 3. Daily sums of measured and modeled sap flow. Two models were used. One (equation 5, model without SWP) was parameterized on the period with the unlimited soil water availability (from 27 February to 5 April, 2008) and these parameters were used to simulate the sap flow during entire season. Second one (equation 9, model including SWP) included soil water potential into the simulation.

Fig. 4. Examples of diurnal curves of measured sap flow (black line), modeled values using the proposed approach (red line), and the classical approach (green line), (panels A, D). Typical diurnal courses of stomatal conductance (panel B, E) estimated using the classical approach from equation 7 (green line) and using the proposed approach (red line). Corresponding global radiation (R_g) and vapor pressure deficit (D) (panel C, F) for days with high (panels A, B, C) and low D (panels D, E, F).

Fig. 5. Measured and modeled values of sap flow using the proposed (A) and the classical approach (B) during the wet period ($n = 955$).

Fig. 6. Response of relative canopy conductance to global radiation (panel A) and canopy conductance to vapor pressure deficit (panel B) during the wet period estimated using the modified Lohammar equation.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6





[Click here to access/download](#)

Electronic Supplementary Material
porovnaní změny June_September.docx

