

NOON state of Bose atoms in the double-well potential via an excited state quantum phase transition

A. A. Bychek^{1,2}, D. N. Maksimov^{1,3}, and A. R. Kolovsky^{1,2}

¹*Kirensky Institute of Physics, Federal Research Center KSC SB RAS, 660036, Krasnoyarsk, Russia*

²*Siberian Federal University, 660041, Krasnoyarsk, Russia*

³*Reshetnev Siberian State University of Science and Technology, 660037, Krasnoyarsk, Russia*

(Dated: June 4, 2018)

We suggest a simple scheme for creating a NOON state of repulsively interacting Bose atoms in the double-well potential. The protocol consists of two steps. First, by setting atom-atom interactions to zero, the system is driven to the upper excited state. Second, the interactions is slowly increased and, simultaneously, the inter-well tunneling is decreased to zero. We analyze fidelity of the final state to the NOON state depending on the number of atoms, ramp rate, and fluctuations of the system parameters. It is shown that for a given fidelity the ramp rate scales algebraically with the number of atoms.

I. INTRODUCTION

Non-classical states of bosonic ensembles play important role in quantum computing, measurement, and communication [1–3]. Among many different implementations [4–10] the two-mode Bose-Hubbard model [11–36] is the most popular playground thanks to its versatility, relative simplicity, and experimental accessibility with ultracold atoms in optical potentials [2, 18, 23, 26, 34, 35]. In this paper we propose a recipe for generating NOON states [37], also known as large cat states [12], in the two-site Bose-Hubbard model. It should be mentioned from the very beginning that, due to decoherence processes inevitably present in a laboratory experiment (particle losses, fluctuations of the optical potential, etc.), the NOON state can be obtained only for relatively small number of atoms. In other words, in the thermodynamic limit $N \rightarrow \infty$ one always gets a statistical mixture of two states with all atoms localized in either of two wells – the phenomenon known as spontaneous localization or parity-symmetry breaking phase transition. One of the goals of this work is to estimate the maximal number of atoms for which one can create the NOON state with the present day experimental facilities.

Formally, the NOON state is the ground state of the one-dimensional *attractive* Bose-Hubbard model [13, 17, 20] in the strong interaction regime. In practice, however, this state is hard to reach because the NOON state are fragile to particle losses caused by the collision instability [38]. To avoid this problem we consider *repulsive* atom-atom interactions where the NOON state appears to be the upper energy state of the system. In what follows we show that this state can be reached in the course of adiabatic passage through an excited state quantum phase transition (ESQPT) [39–41]. It is generally believed that such an adiabatic passage would require extremely long evolution time, which scales exponentially with the number of particles [21]. Here, by detailed examination of the system spectrum in a view of the Landau-Zenner tunnelling, we demonstrate that the evolution time actually scales *algebraically* with the number of bosons N .

A pseudo-classical interpretation of the adiabatic passage with the ESQPT corresponding to a separatrix crossing in the classical phase-space is provided.

II. SYSTEM OVERVIEW

We consider $N \gg 1$ Bose atoms with repulsive interactions in the double-well potential. This system is known to be well described by the two-site Bose-Hubbard Hamiltonian [13, 14]

$$\hat{H} = -\frac{J}{2} (\hat{a}_2^\dagger \hat{a}_1 + \hat{a}_2 \hat{a}_1^\dagger) + \frac{U}{2} \sum_{l=1,2} \hat{n}_l (\hat{n}_l - 1) + \delta (\hat{n}_2 - \hat{n}_1), \quad (1)$$

where J is the hopping matrix element, U the microscopic interaction constant, \hat{a}_l and \hat{a}_l^\dagger the bosonic annihilation and creation operators, \hat{n}_l the number operator, and δ the difference between the on-site energies. For N bosons the Hilbert space of the Hamiltonian (1) of dimension $\mathcal{N} = N + 1$ is spanned by Fock states

$$|N_1, N_2\rangle = |N/2 - n, N/2 + n\rangle \equiv |n\rangle, \quad |n| \leq N/2, \quad (2)$$

where $N_1 + N_2 = N$. Above we used a symmetric parameterization to label the Fock states by a single quantum number n (N is assumed to be even). The full spectrum of the system is shown in Fig.1, where we introduced the macroscopic interaction constant $g = UN/2$ and set the hopping matrix element to $J = 1 - g$. Thus, the case $g = 0$ corresponds to the system of non-interacting bosons while in the case $g = 1$ the inter-well tunneling is completely suppressed. It is easy to prove that the spectrum is equidistant for $g = 0$ and quadratic for $g = 1$, with all energy levels except the ground state being twofold degenerate [22]. The spectrum for intermediate g possesses the quantum separatrix and can be understood by employing the pseudo-classical approach which we review in Sec. III.

Among the eigenstates $|\Psi_j\rangle$ of the Hamiltonian (1) of particular interest are the states with minimal and maximal energy. For $g = 0$ the ground state of the system is




FIG. 1: Energy spectrum of $N = 10$ (left) and $N = 40$ (right) bosons against the macroscopic interaction constant $g = UN/2$. (The other parameters are $J = 1 - g$ and $\delta = 0$.) The quantum separatrix is marked by the red dashed line.

a Bose-Einstein condensate with all particles occupying the symmetric single-particles state,

$$|\Psi_0(g=0)\rangle = \frac{1}{\sqrt{2^N N!}} (a_1^\dagger + a_2^\dagger)^N |vac\rangle, \quad (3)$$

while the upper energy state is a Bose-Einstein condensate with all particles occupying the antisymmetric single-particles state,

$$|\Psi_N(g=0)\rangle = \frac{1}{\sqrt{2^N N!}} (a_1^\dagger - a_2^\dagger)^N |vac\rangle. \quad (4)$$

Let us follow these states under variation of g . At each value of g eigenfunctions are found as an expansion over the Fock states (2),

$$|\Psi_j(g)\rangle = \sum_{n=-N/2}^{N/2} c_n^{(j)}(g) |n\rangle, \quad (5)$$

For $j = 0$ and $j = N$ the results are shown in Fig 2. It is seen that the ground state transforms into the fragmented condensate [22]

$$|\Psi_0(g=1)\rangle = |N/2, N/2\rangle, \quad (6)$$

while the upper energy state evolves into the NOON state

$$|\Psi_N(g=1)\rangle = |NOON\rangle \equiv \frac{1}{\sqrt{2}} (|N, 0\rangle + |0, N\rangle). \quad (7)$$

Next we consider time evolution of the system according to the Schrödinger equation,

$$i \frac{d}{dt} |\psi\rangle = \hat{H}(g) |\psi\rangle, \quad J = 1 - g, \quad (8)$$




FIG. 2: Squared absolute values of expansion coefficients Eq. (5) of the ground (left) and upper energy (right) states against the macroscopic interaction constant g .




FIG. 3: Panel (a): Squared absolute values of expansion coefficients over the Fock basis as the function of $g = \nu t$ for the adiabatic passage with $\nu = 0.1$. Panels (b) and (c) compares final state of the system for $\nu = 0.1$ and $\nu = 0.025$.

with the interaction constant g growing linearly from 0 to 1 during the time interval $T = 1/\nu$. In Fig. 3(a) we present the results of numerical simulations of the system dynamics for $|\psi(t=0)\rangle = |\Psi_N(g=0)\rangle$ and $\nu = 0.1$. Shown are squared absolute values of the expansion coefficients $c_n(t) = \langle n | \psi(t) \rangle$. One can see in Fig. 3(b) that the final state $|\psi(t=T)\rangle$ does not ideally coincide with the target NOON state Eq. (7). With a smaller ramp rate, however, the result is almost perfect, see Fig. 3(c). In the next sections we analyze the discussed adiabatic passage in more detail and quantify the final state $|\psi(t=T)\rangle$. To pay credits to other works we mention that adiabatic pas-

sage for the ground state of the attractive Bose-Hubbard model was considered earlier in Ref. [20, 42] and a different adiabatic passage, which involves the rising potential barrier which separates a Bose-Einstein condensate into two parts, was analyzed in Ref. [14, 15, 19, 24].

III. PSEUDO-CLASSICAL APPROACH

To get a deeper insight into the discussed adiabatic passage we resort to the pseudo-classical approach. This approach borrows its ideas from the semi-classical method in single-particle quantum mechanics to address the spectral and dynamical properties of the system of N interacting bosons, with $1/N$ playing the role of Planck's constant [43–46]. Formally, the creation and annihilation operators are substituted with C -numbers as $\hat{a}_l/\sqrt{N} \rightarrow a_l$ and $\hat{a}_l^\dagger/\sqrt{N} \rightarrow a_l^*$, which also implies rescaling of the Hamiltonian as $\hat{H}/N \rightarrow H$. For the two-site Bose-Hubbard model this leads to the classical Hamiltonian [11]

$$H = \frac{g}{2}I^2 - \frac{J}{2}\sqrt{1-I^2}\cos\phi, \quad J = 1 - g, \quad (9)$$

where $g = UN/2$ is the macroscopic interaction constant.

Fig. 4 shows the phase portrait of the system (9) at four different values of g : 0, 0.4, 0.7, and 1. By using the relation $I = \sin\theta$ phase portraits of the system can be also drawn on sphere of the unit radius. In this representation the line $I = 1$ ($I = -1$) reduces to single point – the north (south) pole of the sphere. For $g = 0$ the phase portrait contains two elliptic points at $(I, \phi) = (0, 0)$ (minimal energy) and $(I, \phi) = (0, \pi)$ (maximal energy), see Fig.4(a). As g is increased above $g_{cr} = 1/3$ the latter elliptic point bifurcates into two elliptic points at $(I, \phi) = (\pm I^*, \pi)$, where I^* is a function of g . With a further increase of




FIG. 4: Phase portraits of the classical Hamiltonian (9) for (a) $g = 0$, (b) $g = 0.4$, (c) $g = 0.7$, and (d) $g = 1$.




FIG. 5: Comparison between the classical (left panel) and quantum (right panel) dynamics. Parameters are $\delta = 0$, $\nu = 0.1$, and $N = 100$.

g the island around the point $(I, \phi) = (0, 0)$ vanishes while the islands around $(I, \phi) = (\pm I^*, \pi)$ monotonically grow, finally leading to the phase-space portrait shown in Fig.4(d).

The depicted phase portraits suffice to find the energy spectrum shown in Fig. 1 by using the semiclassical quantization rule, where the phase volume encircled by a trajectory is required to be a multiple of the effective Planck constant $h = 1/N$. Then the central island around the point $(I, \phi) = (0, 0)$ gives energy levels below the quantum separatrix while two symmetric islands around $(I, \phi) = (\pm I^*, 0)$ give degenerate levels above the quantum separatrix. The details are given in Ref. [45], where it was demonstrated that the pseudo-classical approach provides an accurate approximation to the exact spectrum even for $N = 10$.

Let us now study dynamics of the classical system (9) when both g and J vary in time as $g = \nu t$ and $J = 1 - g$. As the initial condition we take an ensemble of particles with the probability distribution given by the two-dimensional Gaussian centered at the elliptic point $(I, \phi) = (0, \pi)$. For comparison with quantum dynamics the width of the Gaussian is adjusted to $\sigma = \sqrt{N}$. The left panel in Fig. 5 shows the evolution of the classical distribution function $\rho(I, t)$ for $N = 100$. (We stress one more time that the latter parameter determines only the width of the initial distribution.) The left panel in Fig. 5 should be compared with the right panel showing the quantum evolution. The observed agreement underlines the classical phenomenon behind the quantum results discussed in the previous section. Classically, the particles are captured into the upper and lower islands emerging after bifurcation of the elliptic point $(I, \phi) = (0, \pi)$ and then transported towards $I = 1$ and $I = -1$, respectively. The phenomenon of capturing into (and releasing




FIG. 6: Same as in Fig. 5 with $\delta = 0.01$.

from) an elliptic island was considered earlier in Ref. [47] in a different context. It involves the crossing of instantaneous separatrix that, in turn, was analyzed in Ref. [48].

To conclude this section we discuss the effect of non-zero δ . For $\delta \neq 0$ the emerging islands have different size, which makes $\rho(I, t)$ asymmetric with respect to $I \rightarrow -I$. To characterize this asymmetry we introduce the population imbalance

$$G = \int_0^1 \rho(I, T) dI - \int_{-1}^0 \rho(I, T) dI. \quad (10)$$

If δ is increased the population imbalance (10) grows monotonically, approaching $|G| = 1$, see Fig. 6. Importantly, the imbalance also grows if ν is decreased and for any finite δ the imbalance is unity in the limit $\nu \rightarrow 0$.

IV. LANDAU-ZENER TUNNELING AND FIDELITY TO THE NOON STATE

In the previous section we explained the quantum results depicted in Fig. 2 by using the pseudo-classical approach. The quantum-mechanical explanation of these results is based on the notion of Landau-Zener tunneling. Due to this phenomenon several energy levels become populated as we follow the upper most level in Fig. 1 with a finite sweeping rate. This is illustrated in Fig. 7(a) which shows the populations of the instantaneous energy levels $P_j(t)$,

$$P_j(t) = |\langle \Psi_j(g = \nu t) | \psi(t) \rangle|^2, \quad (11)$$

for $\nu = 0.1$ and $N = 30$. Notice that only even levels are populated because of different symmetry of eigenstates of the Hamiltonian (1) with odd and even index j . To quantify the effect of Landau-Zener tunneling we introduce the fidelity

$$F = |\langle NOON | \psi(T) \rangle|^2, \quad (12)$$




FIG. 7: Populations of the instantaneous energy levels for $N = 30$, $\nu = 0.1$, and $\delta = 0$ (upper panel) and $\delta = 0.0001$ (lower panel).

which characterizes how close the final state is to the target NOON state. In the limit $\nu \rightarrow 0$ it is enough to take into account only the nearest high energy level of the same (even) symmetry, which alone determines fidelity of the final state through the celebrated Landau-Zener equation

$$F = 1 - \exp\left(-\frac{\pi\Delta^2}{2|\alpha|\nu}\right). \quad (13)$$

In Eq. (13) Δ is the energy gap between the upper most level and the next level of the same symmetry, $\nu = 1/T$ the sweeping rate, and α is determined by the angle at which two levels approach each other. Since the energy gap Δ and $|\alpha|$ scales algebraically with $1/N$, we expect that the evolution time T has to be increased proportionally to the number of particles to insure a given fidelity. Direct numerical simulations of the adiabatic passage for different N confirm this hypothesis, see Fig. 8(a). It is interesting to discuss the depicted result with respect to the recent laboratory experiment [34] which studies the parity-symmetry-breaking phase transition for $N \approx 4500$ attractively interacting atoms in a double-well potential. Taking $J/h = 40Hz$ and the evolution time $\sim 1s$ we get $N \approx 30$ and this number can be easily increased by one order of magnitude by relaxing the fidelity to $F = 0.9$ and using a time-dependent sweeping rate $\nu = \nu(t)$ that optimizes the adiabatic passage. We stress that the above estimate is obtained under the assumption of negligible decoherence processes which we shall discuss in Sec. V.

Next we analyze the effect of non-zero $\delta \ll J$ in the Hamiltonian (1) from the quantum-mechanical viewpoint. Non-zero δ breaks the reflection symmetry of the system, so that eigenstates of the Hamiltonian (1) at $g \gg J$ are given by the Fock states $|N/2 - n, N/2 + n\rangle$ and $|N/2 + n, N/2 - n\rangle$ but not their symmetric or antisymmetric superpositions. (In particular, $|\Psi_N \approx |N, 0\rangle$ and




FIG. 8: Left panel: Minimal evolution time T insuring fidelity $F = 0.99$ versus the number of bosons N . Right panel: Population imbalance $|G|$ (dashed line) and fidelity F (solid line) as the function of δ for $N = 40$ and $\nu = 0.025$.

$\Psi_{N-1} \approx |0, N\rangle$.) This drastically changes Fig. 7(a) – now both odd and even instantaneous energy levels become populated during the adiabatic passage, see Fig. 7(b). For the considered extremely small value of δ this difference simply reflects a change of the basis and, physically, both Fig. 7(a) and Fig. 7(b) describe the same process, which results in the NOON state as the final state of the system. However, for a larger δ we see considerable deviation from the NOON state, see Fig. 8(b). In particular, in full analogy with the classical result, the population imbalance $|G|$ approaches the unity if $|\delta|$ is increased.

V. DECOHERENCE EFFECTS

The result depicted in Fig. 8(a) proves that, at least in principle, one can create arbitrary large cat state by simply increasing the duration of the adiabatic passage proportionally to the number of particles N . This, however, implicitly assumes the absence of any decoherence process [21] and precision control over the system parameters, in the first place, over parameter δ . In this section we discuss decoherence caused by fluctuations of δ , which are unavoidable in a laboratory experiment.

In the presence of fluctuations the fidelity(12) should be redefined as

$$F = \langle NOON | \mathcal{R}(T) | NOON \rangle, \quad \mathcal{R}(t) = \overline{|\psi(t)\rangle\langle\psi(t)|}, \quad (14)$$

where the bar denotes the average over fluctuations. To be specific, we assume that $\delta(t)$ is the white noise with vanishing mean value, i.e., $\overline{\delta(t)\delta(t')} = \delta_0^2 \delta(t-t')$. Then the density matrix $\mathcal{R}(t)$ is easy to show to obey the following master equation [49]

$$\frac{d\mathcal{R}}{dt} = -i[\hat{H}, \mathcal{R}] - \delta_0^2 [\hat{n}, [\hat{n}, \mathcal{R}]], \quad (15)$$

where $\hat{n} = \hat{n}_1 - \hat{n}_2$. We solve Eq. (15) for the adiabatic passage discussed above. Fig. 9 shows fidelity (14) as the function of the noise amplitude δ_0 for three system sizes $N = 10, 20, 40$, where we proportionally decreased




FIG. 9: Fidelity (14) as the function of the noise amplitude for $N = 10$ and $\nu = 0.1$ (solid line), $N = 20$ and $\nu = 0.05$ (dash-dotted line), and $N = 40$ and $\nu = 0.025$ (dashed line). Insets show the initial density matrix for $N = 20$ (lower-left corner) and final density matrices for $\delta_0 = 0$ and $\delta_0 = 0.1/N$ (upper-right corner).

the sweeping rate ν to insure fidelity $F \approx 1$. One striking feature of the depicted functions is a rapid decay of fidelity to $F \approx 0.5$ in the interval $0 < \delta_0 < \delta_0^*$ where $\delta_0^* = \delta_0^*(N)$. In this interval the off-diagonal elements of the density matrix $\mathcal{R}(T)$ gradually vanish. On the other hand, the diagonal elements of the density matrix remain essentially unaffected. Clearly, this result illustrates the usual quantum-to-classical transition due to a decoherence process [49–51]. Notice that the larger system is, the more it is sensitive to decoherence. Numerical results depicted in Fig. 9 indicate that δ_0^* decreases with N faster than $1/N$.

Next we briefly discuss decoherence due to particle losses. In the case of not conserved number of particles the master equation on the system density matrix reads

$$\frac{d\mathcal{R}}{dt} = -i[\hat{H}, \mathcal{R}] - \gamma \sum_{l=1,2} (\hat{a}_l^\dagger \hat{a}_l \mathcal{R} - 2\hat{a}_l \mathcal{R} \hat{a}_l^\dagger + \mathcal{R} \hat{a}_l^\dagger \hat{a}_l), \quad (16)$$

where γ denotes the decay rate (see recent review Ref. [52]). The value of γ in Eq. (16) crucially depends on the sign of interatomic interactions. For example, in the already cited experiment [35] with attractively interacting atoms the decay rate was $\gamma \approx 0.1$ atoms per second, which make impossible generation of the NOON state even for $N \sim 10$ atoms. On the other hand, it is known that a Bose-Einstein condensate of repulsively interacting atoms may have life-time up to few hours that assumes $\gamma \sim 10^{-4}$. The negligible decoherence rate due to particle losses is our main reason for considering the adiabatic passage for the upper energy state of repulsively interacting atoms instead of that for the ground state of attractively interacting atoms. In all other aspects there is no conceptual difference between the adiabatic passage for the ground and upper states.




FIG. 10: Populations of eigenstates of the Hamiltonian (1) as the function of time. (Note that for $g = 0$ the eigenstates of (1) are given by $|N - n, n\rangle$ where n now denotes the number of particles in the antisymmetric single-particle state.) Parameters are $N = 40$, $g = 0$, $J = 1$, $\omega = J$, $\delta_0 = 0.05$ (left panel) and $N = 40$, $g = 0$, $J = 0.01$, $\delta = 1$ (right panel).

VI. PREPARATION OF THE EXCITED STATE

Finally, we discuss a method to excite the system of non-interacting bosons ($g = 0$) into the highest energy state. A way to do this is to drive the system by periodically changing parameter δ as $\delta(t) = \delta_0 \sin(\omega t)$, where the frequency ω coincides with the transition frequency between the symmetric and antisymmetric single-particle states uniquely determined by the parameter J . If $\delta_0 \ll J$ (the latter condition justifies the rotating-wave approximation) the problem can be solved analytically and leads to the Rabi oscillations, see Fig. 10(a). Thus, to excite the system in the upper state, we need to drive it for one half of the Rabi period.

Another, perhaps even simpler way to obtain the excited state (4) is to quench the system into the parameter region $\delta \gg J$ by suddenly tilting the double-well. Then the time evolution of the expansion coefficients is approximately given by $c_n^{(j)}(t) = \exp(i2\delta nt)c_n^{(j)}(0)$ and after one half of the period $T_B = \pi/\delta$ (which can be interpreted as the Bloch period) the state (3) transforms into the state (4), see Fig. 10(b).

VII. CONCLUSIONS

We suggested a method for creating the NOON state of Bose atoms, i.e., coherent superposition of two states in which all particles are in the same well of the double-well potential. Unlike to previous studies, which almost exclusively focused on the case of attractive interactions [53], we considered the repulsively interacting atoms that avoids the problem of particle losses. The scheme protocol consists of two steps. First, by setting the interatomic interactions to zero we transfer the system from the ground state to the upper excited state. Second, adiabatically increasing the interaction strength and simultaneously decreasing the hopping rate we transform this excited state to the NOON state. In the Fock space the latter stage can be viewed as splitting of the initially localized wave packet into two packets [54]. This process was shown to have a pseudo-classical counterpart and some of quantum results, for example, the population imbalance G can be obtained by using pure classical arguments. Of course, the classical approach cannot address phase coherence between the packets, which is characterized by the fidelity F .

Formally, the suggested scheme allows us to create an arbitrary large cat state. However, any experimental realization of the scheme protocol imposes fundamental limitation on the number of atoms due to decoherence processes present in a laboratory experiment. Here, we analyzed the decoherence caused by fluctuation of the parameter δ (the energy mismatch between the left and right wells of the double-well potential) that appears to be crucial for the system dynamics. It was shown that there is a critical value for the fluctuation amplitude $\delta_0^* \sim 1/N$ above which the final state of the system becomes ‘classical NOON state’, i.e., incoherent superposition of two states in which all particles are in the same well of the double-well potential. Thus to get the NOON state with a large number of atoms every effort to reduce fluctuation of δ should be taken.

Acknowledgements. The authors acknowledge financial support from Russian Foundation for Basic Research, Government of Krasnoyarsk Territory, and Krasnoyarsk Region Science and Technology Support Fund through the grant No. 16-42-240746.

-
- [1] D. J. Wineland, *Nobel Lecture: Superposition, entanglement, and raising Schrödinger’s cat*, *Reviews of Modern Physics* **85**, 1103 (2013).
 - [2] C. Lee, J. Huang, H. Deng, H. Dai, and J. Xu, *Nonlinear quantum interferometry with Bose condensed atoms*, *Frontiers of Physics* **7**, 109 (2012).
 - [3] L. Pezzè, A. Smerzi, M. K. Oberthaler, R. Schmied, and P. Treutlein, *Non-classical states of atomic ensembles: fundamentals and applications in quantum metrology*, arXiv:1609.01609 (2016).

- [4] C. Monroe, D. M. Meekhof, B. E. King, and D. J. Wineland, *A Schrödinger cat superposition state of an atom*, *Science* **272**, 1131 (1996).
- [5] P. C. Haljan, P. J. Lee, K. A. Brickman, M. Acton, L. Deslauriers, and C. Monroe, *Entanglement of trapped-ion clock states*, *Physical Review A* **72**, 062316 (2005).
- [6] M. J. McDonnell, J. P. Home, D. M. Lucas, G. Imreh, B. C. Keitch, D. J. Szwer, N. R. Thomas, S. C. Webster,

- D. N. Stacey, and A. M. Steane, *Long-lived mesoscopic entanglement outside the Lamb-Dicke regime*, Physical Review Letters **98**, 063603 (2007).
- [7] H.-Y. Lo, D. Kienzler, L. de Clercq, M. Marinelli, V. Negnevitsky, B. C. Keitch, and J. P. Home, *Spin-motion entanglement and state diagnosis with squeezed oscillator wavepackets*, Nature **521**, 236 (2015).
- [8] U. R. Fischer and M.-K. Kang, *Photonic cat states from strongly interacting matter waves*, Physical Review Letters **115**, 260404 (2015).
- [9] D. Kienzler, C. Flühmann, V. Negnevitsky, H.-Y. Lo, M. Marinelli, D. Nadlinger, and J. Home, *Observation of quantum interference between separated mechanical oscillator wave packets*, Physical Review Letters **116**, 140402 (2016).
- [10] W.-W. Zhang, S. K. Goyal, F. Gao, B. C. Sanders, and C. Simon, *Creating cat states in one-dimensional quantum walks using delocalized initial states*, New Journal of Physics **18**, 093025 (2016).
- [11] A. Smerzi, S. Fantoni, S. Giovanazzi, and S. R. Shenoy, *Quantum coherent atomic tunneling between two trapped Bose-Einstein condensates*, Physical Review Letters **79**, 4950 (1997).
- [12] J. I. Cirac, M. Lewenstein, K. Mølmer, and P. Zoller, *Quantum superposition states of Bose-Einstein condensates*, Physical Review A **57**, 1208 (1998).
- [13] R. W. Spekkens, and J. E. Sipe, *Spatial fragmentation of a Bose-Einstein condensate in a double-well potential*, Physical Review A **59**, 3868 (1999).
- [14] J. Javanainen, and M. Y. Ivanov, *Splitting a trap containing a Bose-Einstein condensate: Atom number fluctuations*, Physical Review A **60**, 2351 (1999).
- [15] C. Menotti, J. R. Anglin, J. I. Cirac, and P. Zoller, *Dynamic splitting of a Bose-Einstein condensate*, Physical Review A **63**, 023601 (2001).
- [16] A. P. Hines, R. H. McKenzie, and G. J. Milburn, *Entanglement of two-mode Bose-Einstein condensates*, Physical Review A **67**, 013609 (2003).
- [17] T.-L. Ho, and C. Ciobanu, *The Schrödinger cat family in attractive Bose gases*, Journal of low temperature physics **135**, 257 (2004).
- [18] M. Albiez, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler, *Direct observation of tunneling and nonlinear self-trapping in a single bosonic Josephson junction*, Physical Review Letters **95**, 010402 (2005).
- [19] L. Pezzé, L. A. Collins, A. Smerzi, G. P. Berman, and A. R. Bishop, *Sub-shot-noise phase sensitivity with a Bose-Einstein condensate Mach-Zehnder interferometer*, Physical Review A **72**, 043612 (2005).
- [20] Chaohong Lee, *Adiabatic Mach-Zehnder interferometry on a quantized Bose-Josephson junction*, Physical Review Letters **97**, 150402 (2006).
- [21] Y. P. Huang, and M. G. Moore, *Creation, detection, and decoherence of macroscopic quantum superposition states in double-well Bose-Einstein condensates*, Physical Review A **73**, 023606 (2006).
- [22] E. J. Mueller, T.-L. Ho, M. Ueda, and G. Baym, *Fragmentation of Bose-Einstein condensates*, Physical Review A **74**, 033612 (2006).
- [23] R. Gati and M. K. Oberthaler, *A bosonic Josephson junction*, Journal of Physics B: Atomic, Molecular and Optical Physics **40**, R61 (2007).
- [24] A. I. Streltsov, O. E. Alon, and L. S. Cederbaum, *Role of excited states in the splitting of a trapped interacting Bose-Einstein condensate by a time-dependent barrier*, Physical Review Letters **99**, 030402 (2007).
- [25] N. Teichmann, and C. Weiss, *Coherently controlled entanglement generation in a binary Bose-Einstein condensate*, Europhysics Letters **78**, 10009 (2007).
- [26] J. Esteve, C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler, *Squeezing and entanglement in a Bose-Einstein condensate*, Nature **455**, 1216 (2008).
- [27] Y. Khodorkovsky, G. Kurizki, and A. Vardi, *Decoherence and entanglement in a bosonic Josephson junction: Bose-enhanced quantum Zeno control of phase diffusion*, Physical Review A **80**, 023609 (2009).
- [28] T. J. Haigh, A. J. Ferris, and M. K. Olsen, *Demonstrating mesoscopic superpositions in double-well Bose-Einstein condensates*, Optics Communications **283**, 3540 (2010).
- [29] G. Mazzarella, L. Salasnich, A. Parola, and F. Toigo, *Coherence and entanglement in the ground state of a bosonic Josephson junction: From macroscopic Schrödinger cat states to separable Fock states*, Physical Review A **83**, 053607 (2011).
- [30] M. A. Garcia-March, D. R. Dounas-Frazer, and L. D. Carr, *Macroscopic superposition states of ultracold bosons in a double-well potential*, Frontiers of Physics **7**, 131 (2012).
- [31] L. Dell'Anna, *Analytical approach to the two-site Bose-Hubbard model: From Fock states to Schrödinger cat states and entanglement entropy*, Physical Review A **85**, 053608 (2012).
- [32] J. Javanainen, and H. Chen, *Ground state of the double-well condensate for quantum metrology*, Physical Review A **89**, 033613 (2014).
- [33] T. J. Volkoff, *Optimal and near-optimal probe states for quantum metrology of number-conserving two-mode bosonic Hamiltonians*, Physical Review A **94**, 042327 (2016).
- [34] A. Trenkwalder, G. Spagnolli, G. Semeghini, S. Coop, M. Landini, P. Castilho, L. Pezzé, G. Modugno, M. Inguscio, A. Smerzi, and M. Fattori, *Quantum phase transitions with parity-symmetry breaking and hysteresis*, Nature physics **12**, 826 (2016).
- [35] G. Spagnolli, G. Semeghini, L. Masi, G. Ferioli, A. Trenkwalder, S. Coop, M. Landini, L. Pezzé, G. Modugno, M. Inguscio, A. Smerzi, and M. Fattori, *Crossing Over from Attractive to Repulsive Interactions in a Tunneling Bosonic Josephson Junction*, Physical Review Letters **118**, 230403 (2017).
- [36] M. Bilardello, A. Trombettoni, and A. Bassi, *Collapse in ultracold Bose Josephson junctions*, Physical Review A **95**, 032134 (2017).
- [37] H. Lee, P. Kok, and J. P. Dowling, *A quantum Rosetta stone for interferometry*, Journal of Modern Optics **49**, 2325 (2002).
- [38] R. J. Dodd, M. Edwards, C. J. Williams, C. W. Clark, M. J. Holland, P. A. Ruprecht, and K. Burnett, *Role of attractive interactions on Bose-Einstein condensation*, Physical Review A **54**, 661 (1996).
- [39] M. A. Caprio, P. Cejnar, and F. Iachello, *Excited state quantum phase transitions in many-body systems*, Annals of Physics **323**, 1106 (2008).
- [40] P. Pérez-Fernández, A. Relaño, J. M. Arias, P. Cejnar, J. Dukelsky, and J. E. García-Ramos, *Excited-state phase transition and onset of chaos in quantum optical models*, Physical Review E **83**, 046208 (2011).

- [41] A. Relaño, J. Dukelsky, P. Pèrez-Fernàndez, and J. M. Arias, *Quantum phase transitions of atom-molecule Bose mixtures in a double-well potential*, Physical Review E **90**, 042139 (2014).
- [42] Chaohong Lee, *Universality and anomalous mean-field breakdown of symmetry-breaking transitions in a coupled two-component Bose-Einstein condensate*, Physical Review Letters **102**, 070401 (2009).
- [43] K. W. Mahmud, H. Perry, and W. P. Reinhardt, *Quantum phase-space picture of Bose-Einstein condensates in a double well*, Physical Review A **71**, 023615 (2005).
- [44] S. Mossmann, and C. Jung, *Semiclassical approach to Bose-Einstein condensates in a triple well potential*, Physical Review A **74**, 033601 (2006).
- [45] E. M. Graefe, and H. J. Korsch, *Semiclassical quantization of an N -particle Bose-Hubbard model*, Physical Review A **76**, 032116 (2007).
- [46] T. Zibold, E. Nicklas, C. Gross, and M. K. Oberthaler, *Classical bifurcation at the transition from Rabi to Josephson dynamics*, Physical Review Letters **105**, 204101 (2010).
- [47] A. R. Kolovsky, and H. J. Korsch, *Adiabatic scattering of atoms by a standing laser wave*, Physical Review A **55**, 4433 (1997).
- [48] J. H. Hannay, *Accuracy loss of action invariance in adiabatic change of a one-freedom Hamiltonian*, Journal of Physics A: Mathematical and General **19**, L1067 (1986).
- [49] A. R. Kolovsky, *Condition of correspondence between quantum and classical dynamics for a chaotic system* Physical Review Letters **76**, 340 (1996).
- [50] T. Dittrich, and R. Graham, *Effects of weak dissipation on the long-time behaviour of the quantized standard map*, Europhysics Letters **7**, 287 (1988).
- [51] W. H. Zurek, *Decoherence and the transition from Quantum to Classical*, Physics Today **44**, 36 (1991).
- [52] G. Kordas, D. Witthaut, P. Buonsante, A. Vezzani, R. Burioni, A.I. Karanikas, and S. Wimberger, *The dissipative Bose-Hubbard model: Methods and examples*, Eur. Phys. J. Special Topics **224**, 2127 (2015).
- [53] It is appropriate place here to mention the other method for creating the NOON state of repulsively interacting atoms based on the quench [54] or lossy dynamics [55] and the works [?] which explore possibility of using the two-component BEC in a single well/trap instead of the single component BEC in a double-well potential.
- [54] A. I. Streltsov, O. E. Alon, and L. S. Cederbaum, *Efficient generation and properties of mesoscopic quantum superposition states in an attractive Bose-Einstein condensate threaded by a potential barrier*, Journal of Physics B: Atomic, Molecular and Optical Physics **42**, 091004 (2009).
- [55] G. Kordas, S. Wimberger, and D. Witthaut, *Dissipation-induced macroscopic entanglement in an open optical lattice*, Europhysics Letters, **100**, 30007 (2012).
- [56] A. Micheli, D. Jaksch, J. I. Cirac, and P. Zoller, *Many-particle entanglement in two-component Bose-Einstein condensates*, Phys. Rev. A **67**, 013607 (2003).
- [57] L. Pezzè and A. Smerzi, *Entanglement, Nonlinear Dynamics, and the Heisenberg Limit*, Phys. Rev. Lett. **102**, 100401 (2009).
- [58] C. Gross, T. Zibold, E. Nicklas, J. Estève, and M. K. Oberthaler, *Nonlinear atom interferometer surpasses classical precision limit*, Nature **464**, 1165 (2010).
- [59] H. Strobel, W. Muessel, D. Linnemann, T. Zibold, D. B. Hume, L. Pezzè, A. Smerzi, and M. K. Oberthaler, *Fisher information and entanglement of non-Gaussian spin states*, Science **345**, 424 (2014).