

DOI: 10.17516/1997-1397-2020-13-6-755-762

УДК 517.95

Mixed Biharmonic Dirichlet-Neumann Problem in Exterior Domains

Hovik A. Matevossian*

Federal Research Center "Computer Science and Control" RAS
Moscow, Russian Federation
Moscow Aviation Institute (National Research University)
Moscow, Russian Federation

Received 17.09.2019, received in revised form 04.06.2020, accepted 17.10.2020

Abstract. We study the unique solvability of the mixed Dirichlet-Neumann problem for the biharmonic equation in the exterior of a compact set under the assumption that solutions of this problem have bounded Dirichlet integrals with the weight $|x|^a$. Depending on the value of the parameter a , we obtained uniqueness (non-uniqueness) theorems of the problem and present exact formulas for the dimension of the space of solutions of the mixed Dirichlet-Neumann problem.

Keywords: biharmonic operator, Dirichlet-Neumann problem, weighted Dirichlet integral.

Citation: H.A. Matevossian, Mixed Biharmonic Dirichlet-Neumann Problem in Exterior Domains, J. Sib. Fed. Univ. Math. Phys., 2020, 13(6), 755–762. DOI: 10.17516/1997-1397-2020-13-6-755-762.

1. Introduction and preliminaries

Let Ω be an unbounded domain in \mathbb{R}^n , $n \geq 2$, $\Omega = \mathbb{R}^n \setminus \overline{G}$ with the boundary $\partial\Omega \in C^2$, where G is a bounded simply connected domain (or a union of finitely many such domains) in \mathbb{R}^n , $0 \in G$, $\overline{\Omega} = \Omega \cup \partial\Omega$ is the closure of Ω , $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $|x| = \sqrt{x_1^2 + \dots + x_n^2}$.

In the domain Ω we consider the following mixed problems for the biharmonic equation

$$\Delta^2 u = 0 \tag{1}$$

with the Dirichlet–Neumann boundary conditions

$$u|_{\Gamma_1} = \frac{\partial u}{\partial \nu}|_{\Gamma_1} = 0, \quad \Delta u|_{\Gamma_2} = \frac{\partial \Delta u}{\partial \nu}|_{\Gamma_2} = 0, \tag{2}$$

where $\overline{\Gamma_1} \cup \overline{\Gamma_2} = \partial\Omega$, $\Gamma_1 \cap \Gamma_2 = \emptyset$, $\text{mes}_{n-1} \Gamma_1 \neq 0$, $\nu = (\nu_1, \dots, \nu_n)$ is the outer unit normal vector to $\partial\Omega$.

As is well known, if Ω is an unbounded domain, one should additionally characterize the behavior of the solution at infinity. As a rule, to this end, one usually poses either the condition that the Dirichlet (energy) integral is finite or a condition on the character of vanishing of the modulus of the solution as $|x| \rightarrow \infty$. Such conditions at infinity are natural and were studied by several authors (e.g., [6–8]).

Elliptic problems with parameters in the boundary conditions have been called Steklov or Steklov-type problems, since their first appearance in [27]. For the biharmonic operator, these

*hmatevossian@graduate.org <https://orcid.org/0000-0002-9895-9628>

© Siberian Federal University. All rights reserved

conditions were first considered in [1,9] and [25], where the isoperimetric properties of the first eigenvalue were studied.

Note that standard elliptic regularity results are available in [3]. The monograph covers higher order linear and nonlinear elliptic boundary value problems, mainly with the biharmonic or polyharmonic operator as the leading principal part. The underlying models and, in particular, the role of different boundary conditions are explained in detail. As for linear problems, after a brief summary of the existence theory and L^p and Schauder estimates, the focus is on positivity. The required kernel estimates are also presented in detail.

In [2], the boundary value problems for the biharmonic equation and the Stokes system are studied in a half space, and, using the Schwarz reflection principle in weighted L^q -space, the uniqueness of solutions of the Stokes system or the biharmonic equation is proved.

In the present note, this condition is the boundedness of the weighted Dirichlet integral:

$$D_a(u, \Omega) \equiv \int_{\Omega} |x|^a \sum_{|\alpha|=2} |\partial^\alpha u|^2 dx < \infty, \quad a \in \mathbb{R}.$$

In various classes of unbounded domains with finite weighted Dirichlet (energy) integral, one of the author [10–23] studied uniqueness (non–uniqueness) problem and found the dimensions of the spaces of solutions of boundary value problems for the elasticity system and the biharmonic (polyharmonic) equation.

By developing an approach based on the use of Hardy type inequalities [6–8], in the present note, we obtain a uniqueness (non–uniqueness) criterion for a solution of the mixed Dirichlet–Neumann problem for the biharmonic equation.

Notation: $C_0^\infty(\Omega)$ is the space of infinitely differentiable functions in Ω with compact support in Ω . We denote by $H^m(\Omega, \Gamma)$, $\Gamma \subset \bar{\Omega}$, the Sobolev space of functions in Ω obtained by the completion of $C^\infty(\bar{\Omega})$ vanishing in a neighborhood of Γ with respect to the norm

$$\|u; H^m(\Omega, \Gamma)\| = \left(\int_{\Omega} \sum_{|\alpha| \leq m} |\partial^\alpha u|^2 dx \right)^{1/2}, \quad m = 1, 2,$$

where $\partial^\alpha \equiv \partial^{|\alpha|} / \partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}$, $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multi-index, $\alpha_i \geq 0$ are integers, and $|\alpha| = \alpha_1 + \dots + \alpha_n$; if $\Gamma = \emptyset$, we denote $H^m(\Omega, \Gamma)$ by $H^m(\Omega)$.

$\overset{\circ}{H}{}^m(\Omega)$ is the space obtained by the completion of $C_0^\infty(\Omega)$ with respect to the norm $\|u; \overset{\circ}{H}{}^m(\Omega)\|$.

$\overset{\circ}{H}_{loc}{}^m(\Omega)$ is the space obtained by the completion of $C_0^\infty(\Omega)$ with respect to the family of semi-norms

$$\|u; H^m(\Omega \cap B_0(R))\| = \left(\int_{\Omega \cap B_0(R)} \sum_{|\alpha| \leq m} |\partial^\alpha u|^2 dx \right)^{1/2}$$

for all open balls $B_0(R) := \{x : |x| < R\}$ in \mathbb{R}^n for which $\Omega \cap B_0(R) \neq \emptyset$.

Let $\binom{n}{k}$ be the (n, k) -binomial coefficient, $\binom{n}{k} = 0$ for $k > n$.

2. Definitions and auxiliary statements

Definition 2.1. A solution of the homogenous biharmonic equation (1) in Ω is a function $u \in H_{loc}^2(\Omega)$ such that for every function $\varphi \in C_0^\infty(\Omega)$, the following integral identity holds:

$$\int_{\Omega} \Delta u \Delta \varphi dx = 0.$$

Lemma 2.2. *Let u be a solution of equation (1) in Ω such that $D_a(u, \Omega) < \infty$. Then*

$$u(x) = P(x) + \sum_{\beta_0 < |\alpha| \leq \beta} \partial^\alpha \Gamma(x) C_\alpha + u^\beta(x), \quad x \in \Omega, \quad (3)$$

where $P(x)$ is a polynomial, $\text{ord } P(x) < m_0 = \max\{2, 2 - n/2 - a/2\}$, $\beta_0 = 2 - n/2 + a/2$, $\Gamma(x)$ is the fundamental solution of equation (1), $C_\alpha = \text{const}$, $\beta \geq 0$ is an integer, and the function u^β satisfies the estimate:

$$|\partial^\gamma u^\beta(x)| \leq C_{\gamma\beta} |x|^{3-n-\beta-|\gamma|}, \quad C_{\gamma\beta} = \text{const},$$

for every multi-index γ .

Remark 2.3. *As is known [26], the fundamental solution $\Gamma(x)$ of the biharmonic equation has the form*

$$\Gamma(x) = \begin{cases} C|x|^{4-n} & \text{if } 4 - n < 0 \text{ or } n \text{ is odd,} \\ C|x|^{4-n} \ln|x| & \text{if } 4 - n \geq 0 \text{ and } n \text{ is even.} \end{cases}$$

Proof of Lemma 2.2. Consider the function $v(x) = \theta_N(x)u(x)$, where $\theta_N(x) = \theta(|x|/N)$, $\theta \in C^\infty(\mathbb{R}^n)$, $0 \leq \theta \leq 1$, $\theta(s) = 0$ for $s \leq 1$, $\theta(s) = 1$ for $s \geq 2$, while $N \gg 1$ and $G \subset \{x : |x| < N\}$. We extend v to \mathbb{R}^n by setting $v = 0$ on $G = \mathbb{R}^n \setminus \bar{\Omega}$.

Then the function v belongs to $C^\infty(\mathbb{R}^n)$ and satisfies the equation

$$\Delta^2 v = f,$$

where $f \in C_0^\infty(\mathbb{R}^n)$ and $\text{supp } f \subset \{x : |x| < 2N\}$. It is easy to see that $D_a(v, \mathbb{R}^n) < \infty$.

We can now use Theorem 1 of [5] since it is based on Lemma 2 of [5], which imposes no constraint on the sign of σ . Hence, the expansion

$$v(x) = P(x) + \sum_{\beta_0 < |\alpha| \leq \beta} \partial^\alpha \Gamma(x) C_\alpha + v^\beta(x),$$

holds for each a , where $P(x)$ is a polynomial of order $\text{ord } P(x) < m_0 = \max\{2, 2 - n/2 - a/2\}$, $\beta_0 = 2 - n/2 + a/2$, $C_\alpha = \text{const}$ and

$$|\partial^\gamma v^\beta(x)| \leq C_{\gamma\beta} |x|^{3-n-\beta-|\gamma|}, \quad C_{\gamma\beta} = \text{const}.$$

Therefore, by the definition of v , we obtain (3). The proof of Lemma 2.2 is complete. \square

Definition 2.4. *A function u is a solution of the mixed Dirichlet–Neumann problem (1), (2), if $u \in \overset{\circ}{H}_{loc}^2(\Omega, \Gamma_1)$ such that for every function $\varphi \in C_0^\infty(\mathbb{R}^n)$, $\varphi = 0$ in the neighborhood of Γ_1 , the following integral identity holds:*

$$\int_{\Omega} \Delta u \Delta \varphi \, dx = 0. \quad (4)$$

3. Main Results

Theorem 3.1. *The mixed Dirichlet–Neumann problem (1), (2) with the condition $D(u, \Omega) < \infty$ has $n + 1$ linearly independent solutions.*

Proof. For any nonzero vector A in \mathbb{R}^n , we construct a generalized solution u_A of the biharmonic equation (1) with the boundary conditions

$$u_A(x)|_{\Gamma_1} = (Ax)|_{\Gamma_1}, \quad \frac{\partial u_A(x)}{\partial \nu}|_{\Gamma_1} = \frac{\partial (Ax)}{\partial \nu}|_{\Gamma_1}, \quad \Delta u_A|_{\Gamma_2} = \frac{\partial \Delta u_A}{\partial \nu}|_{\Gamma_2} = 0, \quad (5)$$

and the condition

$$\chi(u_A, \Omega) \equiv \begin{cases} \int_{\Omega} \left(\frac{|u_A(x)|^2}{|x|^4} + \frac{|\nabla u_A(x)|^2}{|x|^2} + |\nabla \nabla u_A(x)|^2 \right) dx < \infty & \text{for } n > 4, \\ \int_{\Omega} \left(\frac{|u_A(x)|^2}{||x|^2 \ln |x||^2} + \frac{|\nabla u_A(x)|^2}{||x| \ln |x||^2} + |\nabla \nabla u_A(x)|^2 \right) dx < \infty & \text{for } 2 \leq n \leq 4, \end{cases} \quad (6)$$

for $A, x \in \mathbb{R}^n$, where Ax denotes the standard scalar product of A and x .

Such a solution of problem (1), (5) can be constructed by the variational method [26], minimizing the functional

$$\Phi(v) = \frac{1}{2} \int_{\Omega} |\Delta v|^2 dx$$

in the class of admissible functions $\{v: v \in H^2(\Omega), v(x)|_{\Gamma_1} = (Ax)|_{\Gamma_1}, \frac{\partial v(x)}{\partial \nu}|_{\Gamma_1} = \frac{\partial (Ax)}{\partial \nu}|_{\Gamma_1}, v \text{ is compactly supported in } \overline{\Omega}\}$. The validity of condition (6) as a consequence of the Hardy inequality follows from the results in [6–8].

Now, for any arbitrary number $e \neq 0$, we construct a generalized solution u_e of equation (1) with the boundary conditions

$$u_e|_{\Gamma_1} = e, \quad \frac{\partial u_e}{\partial \nu}|_{\Gamma_1} = 0, \quad \Delta u_e|_{\Gamma_2} = \frac{\partial \Delta u_e}{\partial \nu}|_{\Gamma_2} = 0, \quad (7)$$

and the condition

$$\chi(u_e, \Omega) \equiv \begin{cases} \int_{\Omega} \left(\frac{|u_e(x)|^2}{|x|^4} + \frac{|\nabla u_e(x)|^2}{|x|^2} + |\nabla \nabla u_e(x)|^2 \right) dx < \infty & \text{for } n > 4, \\ \int_{\Omega} \left(\frac{|u_e(x)|^2}{||x|^2 \ln |x||^2} + \frac{|\nabla u_e(x)|^2}{||x| \ln |x||^2} + |\nabla \nabla u_e(x)|^2 \right) dx < \infty & \text{for } 2 \leq n \leq 4. \end{cases} \quad (8)$$

The solution of problem (1), (7) is also constructed by the variational method with the minimization of the corresponding functional in the class of admissible functions $\{v : v \in H^2(\Omega), v|_{\Gamma_1} = e, \frac{\partial v}{\partial \nu}|_{\Gamma_1} = 0, v \text{ is compactly supported in } \overline{\Omega}\}$. The condition (8) as a consequence of the Hardy inequality follows from the results in [6–8].

Consider the function $v(x) = (u_A(x) - Ax) - (u_e - e)$. Obviously, v is a solution of problem (1), (2):

$$\Delta^2 v = 0, \quad x \in \Omega, \quad v|_{\Gamma_1} = \frac{\partial v}{\partial \nu}|_{\Gamma_1} = 0, \quad \Delta v|_{\Gamma_2} = \frac{\partial \Delta v}{\partial \nu}|_{\Gamma_2} = 0.$$

One can easily see that $v \not\equiv 0$ and $D(v, \Omega) < \infty$.

To each nonzero vector $\mathbf{A} = (A_0, A_1, \dots, A_n)$ in \mathbb{R}^{n+1} , there corresponds a nonzero solution $v_{\mathbf{A}} = (v_{A_0}, v_{A_1}, \dots, v_{A_n})$ of problem (1), (2) with the condition $D(v_{\mathbf{A}}, \Omega) < \infty$, and moreover,

$$v_{\mathbf{A}}(x) = u_A(x) - u_e - Ax + e.$$

Let A_0, A_1, \dots, A_n be a basis in \mathbb{R}^{n+1} . Let us prove that the corresponding solutions $v_{A_0}, v_{A_1}, \dots, v_{A_n}$ are linearly independent. Let

$$\sum_{i=0}^n C_i v_{A_i} \equiv 0, \quad C_i = \text{const}.$$

Set $W(x) \equiv \sum_{i=1}^n C_i A_i x - C_0 e$. We have $W(x) = \sum_{i=1}^n C_i u_{A_i}(x) - C_0 u_e$,

$$\int_{\Omega} |x|^{-2} |\nabla W|^2 dx < \infty, \quad n > 4; \quad \int_{\Omega} \|x\| \ln \|x\|^{-2} |\nabla W|^2 dx < \infty, \quad 2 \leq n \leq 4.$$

Let us show that

$$W(x) \equiv \sum_{i=1}^n C_i A_i x - C_0 e \equiv 0.$$

Let $T = \sum_{i=0}^n C_i A_i = (t_0, \dots, t_n)$, where $A_0 = -e$. Then

$$\begin{aligned} \int_{\Omega} |x|^{-2} |\nabla W|^2 dx &= \int_{\Omega} |x|^{-2} (t_1^2 + \dots + t_n^2) dx = \infty, \quad n > 4, \\ \int_{\Omega} \|x\| \ln \|x\|^{-2} |\nabla W|^2 dx &= \int_{\Omega} \|x\| \ln \|x\|^{-2} (t_1^2 + \dots + t_n^2) dx = \infty, \quad 2 \leq n \leq 4, \end{aligned}$$

if $T \neq 0$.

Consequently, $T = \sum_{i=0}^n C_i A_i = 0$, and since the vectors A_0, A_1, \dots, A_n are linearly independent, we obtain $C_i = 0, i = 0, 1, \dots, n$.

Thus, the Dirichlet-Neumann problem (1), (2) with the condition $D(u, \Omega) < \infty$ has at least $n + 1$ linearly independent solutions.

Let us prove that each solution u of problem (1), (2) with the condition $D(u, \Omega) < \infty$ can be represented as a linear combination of the functions $v_{A_0}, v_{A_1}, \dots, v_{A_n}$, i.e.

$$u = \sum_{i=0}^n C_i v_{A_i}, \quad C_i = \text{const}.$$

Since A_0, A_1, \dots, A_n is a basis in \mathbb{R}^{n+1} , it follows that there exist constants C_0, C_1, \dots, C_n such that

$$A = \sum_{i=0}^n C_i A_i.$$

We set

$$u_0 \equiv u - \sum_{i=0}^n C_i v_{A_i}.$$

Obviously, the function u_0 is a solution of problem (1), (2), and $D(u_0, \Omega) < \infty, \chi(u_0, \Omega) < \infty$.

Let us show that $u_0 \equiv 0, x \in \Omega$. To this end, we substitute the function $\varphi(x) = u_0(x)\theta_N(x)$ into the integral identity (4) for the function u_0 , where $\theta_N(x) = \theta(|x|/N), \theta \in C^\infty(\mathbb{R}), 0 \leq \theta \leq 1, \theta(s) = 0$ for $s \geq 2$ and $\theta(s) = 1$ for $s \leq 1$; then we obtain

$$\int_{\Omega} (\Delta u_0)^2 \theta_N(x) dx = -J_1(u_0) - J_2(u_0), \tag{9}$$

where

$$J_1(u_0) = 2 \int_{\Omega} \Delta u_0 \nabla u_0 \nabla \theta_N(x) dx, \quad J_2(u_0) = \int_{\Omega} u_0 \Delta u_0 \Delta \theta_N(x) dx.$$

By applying the Cauchy–Schwarz inequality and by taking into account the conditions $D(u_0, \Omega) < \infty$ and $\chi(u_0, \Omega) < \infty$, one can easily show that $J_1(u_0) \rightarrow 0$ and $J_2(u_0) \rightarrow 0$ as $N \rightarrow \infty$. Consequently, by passing to the limit as $N \rightarrow \infty$ in (9), we obtain

$$\int_{\Omega} (\Delta u_0)^2 dx = 0.$$

Therefore, we have

$$\begin{aligned} \Delta u_0 &= 0, \quad x \in \Omega, \\ u_0|_{\Gamma_1} &= \frac{\partial u_0}{\partial \nu} \Big|_{\Gamma_1} = 0, \quad \Delta u_0|_{\Gamma_2} = \frac{\partial \Delta u_0}{\partial \nu} \Big|_{\Gamma_2} = 0. \end{aligned}$$

Hence, it follows [4, Ch.2] that $u_0 \equiv 0$ in Ω . The proof of the theorem is complete. □

Theorem 3.2. *The mixed Dirichlet–Neumann problem (1), (2) with the condition $D_a(u, \Omega) < \infty$ has:*

- (i) *the trivial solution for $n - 2 \leq a < \infty, n > 4$;*
- (ii) *n linearly independent solutions for $n - 4 \leq a < n - 2, n > 4$;*
- (iii) *$n + 1$ linearly independent solutions for $-n \leq a < n - 4, n > 4$;*
- (iv) *$k(r, n)$ linearly independent solutions for $-2r + 2 - n \leq a < -2r + 4 - n, r > 1, n > 4$,*
where

$$k(r, n) = \binom{r+n}{n} - \binom{r+n-4}{n}.$$

The proof of Theorem 3.2 is based on Lemma 2.2 about the asymptotic expansion of the solution of the biharmonic equation and the Hardy type inequalities for unbounded domains [6–8]. In case (iv), we need to determine the number of linearly independent solutions of the biharmonic equation (1), the degree of which do not exceed the fixed number.

It is well know that the dimension of the space of all polynomials in \mathbb{R}^n of degree $\leq r$ is equal $\binom{r+n}{n}$ [24]. Then the dimension of the space of all biharmonic polynomials in \mathbb{R}^n of degree $\leq r$ is equal to

$$\binom{r+n}{n} - \binom{r+n-4}{n},$$

since the biharmonic equation is the vanishing of some polynomial of degree $r - 4$ in \mathbb{R}^n . If we denote by $k(r, n)$ the number of linearly independent polynomial solutions of equation (1) whose degree do not exceed r and by $l(r, n)$ the number of linearly independent homogeneous polynomials of degree r , that are solutions of equation (1), then

$$k(r, n) = \sum_{s=0}^r l(s, n), \quad \text{where } l(s, n) = \binom{s+n-1}{n-1} - \binom{s+n-5}{n-1}, \quad s > 0.$$

Further, we prove that the mixed Dirichlet–Neumann problem (1), (2) with the condition $D_a(u, \Omega) < \infty$ for $-2r + 2 - n \leq a < -2r + 4 - n$ has equally $k(r, n)$ linearly independent solutions.

References

[1] F.Brock, An isoperimetric inequality for eigenvalues of the Stekloff problem, *Z. Angew. Math. Mech. (ZAMM)*, **81**(2001), no. 1, 69–71.

- [2] R.Farwig, A note on the reflection principle for the biharmonic equation and the Stokes system, *Acta Appl. Math.*, **34**(1994), 41–51.
- [3] F.Gazzola, H.-Ch.Grunau, G.Sweers, Polyharmonic Boundary Value Problems: Positivity Preserving and Nonlinear Higher Order Elliptic Equations in Bounded Domains, *Lecture Notes Math.*, vol. 1991, Springer-Verlag, 2010.
- [4] D.Gilbarg, N.Trudinger, *Elliptic Partial Differential Equations of Second Order*. Berlin, Springer-Verlag, 1977.
- [5] V.A.Kondratiev, O.A.Oleinik, On the behavior at infinity of solutions of elliptic systems with a finite energy integral, *Arch. Rational Mech. Anal.*, **99**(1987), no. 1, 75–99.
- [6] V.A.Kondratiev, O.A.Oleinik, Boundary value problems for the system of elasticity theory in unbounded domains. Korn’s inequalities, *Russian Math. Surveys*, **43**(1988), no. 5, 65–119.
- [7] V.A.Kondratiev, O.A.Oleinik, Hardy’s and Korn’s Inequality and their Application, *Rend. Mat. Appl.*, Serie VII, **10**(1990), no. 3, 641–666.
- [8] A.A.Kon’kov, On the dimension of the solution space of elliptic systems in unbounded domains, *Russian Acad. Sci. Sbornik Math.*, **80**(1995), no. 2, 411–434.
- [9] J.R.Kuttler, V.G.Sigillito, Inequalities for membrane and Stekloff eigenvalues, *J. Math. Anal. Appl.*, **23**(1968), no. 1, 148–160.
- [10] O.A.Matevosyan, On solutions of boundary value problems for a system in the theory of elasticity and for the biharmonic equation in a half-space, *Diff. Equations.*, **34**(1998), no. 6, 803–808.
- [11] O.A.Matevosyan, The exterior Dirichlet problem for the biharmonic equation: Solutions with bounded Dirichlet integral, *Math. Notes*, **70**(2001), no. 3, 363–377.
- [12] O.A.Matevosyan, Solutions of exterior boundary value problems for the elasticity system in weighted spaces, *Sbornik Math.*, **192**(2001), no. 12, 1763–1798.
- [13] O.A.Matevosyan, On solutions of mixed boundary-value problems for the elasticity system in unbounded domains, *Izvestiya Math.*, **67**(2003), no. 5, 895–929.
DOI: 10.1070/IM2003v067n05ABEH000451
- [14] O.A.Matevosyan, On solutions of the Dirichlet problem for the polyharmonic equation in unbounded domains, *P-Adic Numbers, Ultrametric Analysis, and Appl.*, **7**(2015), no. 1, 74–78. DOI: 10.1134/S2070046615010069
- [15] O.A.Matevosyan, Solution of a mixed boundary value problem for the biharmonic equation with finite weighted Dirichlet integral, *Diff. Equations*, **51**(2015), no. 4, 487–501.
DOI: 10.1134/S0012266115040060
- [16] O.A.Matevosyan, On solutions of the Neumann problem for the biharmonic equation in unbounded domains, *Math. Notes*, **98**(2015), 990–994. DOI: 10.1134/S0001434615110334
- [17] O.A.Matevosyan, On solutions of the mixed Dirichlet–Navier problem for the polyharmonic equation in exterior domains, *Russ. J. Math. Phys.*, **23**(2016), no. 1, 135–138.
DOI: 10.1134/S106192081601012X

- [18] O.A.Matevosyan, On solutions of one boundary value problem for the biharmonic equation, *Diff. Equations*, **52**(2016), no. 10, 1379–1383. DOI: 10.1134/S0012266116100153
- [19] H.A.Matevossian, On the biharmonic Steklov problem in weighted spaces, *Russ. J. Math. Phys.*, **24**(2017), no. 1, 134–138. DOI: 10.1134/S1061920817010125
- [20] H.A.Matevossian, On solutions of the mixed Dirichlet–Steklov problem for the biharmonic equation in exterior domains, *P-Adic Numbers, Ultrametric Analysis, and Appl.*, **9**(2017), no. 2, 151–157. DOI: 10.1134/S2070046617020054
- [21] H.A.Matevossian, On the Steklov–type biharmonic problem in unbounded domains, *Russ. J. Math. Phys.*, **25**(2018), no. 2, 271–276. DOI: 10.1134/S1061920818020115
- [22] O.A.Matevossian, Mixed Dirichlet–Steklov problem for the biharmonic equation in weighted spaces, *J. Math. Sci. (N. Y.)*, **234**(2018), no 4, 440–454. DOI: 10.1007/s10958-018-4021-8
- [23] H.A.Matevossian, On the polyharmonic Neumann problem in weighted spaces, *Complex Variables and Elliptic Equations*, **64**(2019), no. 1, 1–7. DOI: 10.1080/17476933.2017.1409740
- [24] S.G.Mikhlin, Linear Partial Differential Equations, Vyssaya Shkola, Moscow, 1977 (in Russian).
- [25] L.E.Payne, Some isoperimetric inequalities for harmonic functions, *SIAM J. Math. Anal.*, **1**(1970), no. 3, 354–359.
- [26] S.L.Sobolev, Applications of Functional Analysis in Mathematical Physics, Amer. Math. Soc., Providence, 1991.
- [27] W.Stekloff, Sur les problemes fondamentaux de la physique mathematique, *Ann. Sci. de l'E.N.S.*, 3^e serie, **19**(1902), 191–259 et 455–490.

Смешанная бигармоническая задача Дирихле–Неймана во внешних областях

Овик А. Матевосян

Федеральный исследовательский центр «Информатика и управление» РАН

Москва, Российская Федерация

Московский авиационный институт (национальный исследовательский университет)

Москва, Российская Федерация

Аннотация. Изучаются вопросы единственности решения смешанной задачи Дирихле–Неймана для бигармонического уравнения во внешности компактного множества, в предположении, что обобщенное решение этой задачи обладает конечным интегралом Дирихле с весом $|x|^a$. В зависимости от значения параметра a доказаны теоремы единственности (неединственности), и найдены точные формулы для вычисления размерности пространства решений смешанной задачи Дирихле–Неймана.

Ключевые слова: бигармонический оператор, задача Дирихле–Неймана, весовой интеграл Дирихле.